INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Docetaxel + cyclophosphamide (TC)

Understand your treatment. Know what to expect. Keep your plan to hand.

Displayed calendar
21 days
See the active treatment phase

Treatment stages
1
Explained in the treatment reference

Regimen source
NCCP 00250
Version 9

> Docetaxel and cyclophosphamide on Day 1 every 21 days, for four to six cycles. Confirm your planned number.

Treatment stages

01 **TC**
21-day interval

Your team confirms the active stage and dates. The full treatment guide explains each step.

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	4
03	Symptoms and supportive care	10
04	Sources and further information	15

My name _____

Hospital / oncology team _____

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

Understanding your treatment

Docetaxel + cyclophosphamide (TC) · National Cancer Control Programme (NCCP) 00250

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Report new breathing symptoms, severe diarrhoea, inability to drink, painful swollen limbs, unusual bleeding, or worsening numbness. Tell staff immediately about infusion-site pain or an allergic reaction.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

How the medicines work

Treatment after surgery to reduce the chance of breast cancer returning.

TAXANE CHEMOTHERAPY

Docetaxel



Disrupts the cell’s division machinery

Docetaxel stabilises tiny structures called microtubules. Cells need these to separate their chromosomes during division; disrupting the process can stop cancer cells multiplying. Healthy dividing cells are also affected.


Microtubules stabilised → Division disrupted → Cell growth affected

What to watch for

Take prescribed steroid premedication. Report fever, lasting numbness, swelling, sore nails or infusion symptoms.

ALKYLATING CHEMOTHERAPY

Cyclophosphamide



Damages deoxyribonucleic acid (the cell's genetic material) (DNA) needed for growth

Cyclophosphamide is activated in the body and damages DNA. This helps stop dividing cancer cells but can also affect blood-cell production and reproductive tissues.

Drug activated → DNA damage → Cell division affected

What to watch for

Use the prescribed anti-sickness and hydration plan. Report fever, painful urination or blood in the urine.

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Why this treatment?

TC is an anthracycline-free chemotherapy option after surgery. The team weighs recurrence risk and cancer biology against benefits and side effects. Avoiding an anthracycline can matter for some people, but TC still needs blood tests, infection precautions and a docetaxel steroid plan.

Your treatment setting

Adjuvant high-risk node-positive or node-negative operable breast cancer

The sequence

Docetaxel and cyclophosphamide on Day 1 every 21 days for four to six cycles. The team confirms the planned course before treatment. Radiotherapy, when indicated, follows chemotherapy.

Alternatives and your priorities

Ask what alternatives fit your cancer, what the treatment aims to achieve, and how its practical demands fit your life. The team can discuss personal benefits and risks.

Why this treatment may be chosen

Show trial results and what the numbers mean

- Pembrolizumab + chemotherapy
- Placebo + chemotherapy

Your team can discuss expected benefit and alternatives in the context of your own cancer and priorities.

SACTCHECK · CHAPTER 02

Your schedule, tests and appointments

Docetaxel + cyclophosphamide (TC) · NCCP 00250

The phases and how they connect

TC

21-day view



Chemotherapy · Day 1
Day 1

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

21-DAY INTERVAL

TC

4–6 cycles

Docetaxel and cyclophosphamide on Day 1 every 21 days for four to six cycles. The team confirms the planned course before treatment. Radiotherapy, when indicated, follows chemotherapy.

The treatment phase determines the medicines, supportive prescription and tests. A phase change does not reset previous drug exposure or cardiac monitoring.

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Relative milestones, not equal time intervals. The clinical team owns booking, result review and action on overdue tests. This is an educational map, not an automated appointment or result tracker.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00250 v9 · official regimen · ESMO 2024 early breast cancer guideline · treatment and follow-up · ESC 2022 cardio-oncology guideline · cardiac surveillance and risk · NICE NG101 · bone health and follow-up · ESMO metastatic breast cancer guideline · 2026

THE WHOLE TREATMENT JOURNEY

Docetaxel + cyclophosphamide (TC): treatment and checks

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Early cycles	Later cycles / transition	Next treatment / completion	Follow-up
Cancer medicines NCCP 00250	Agree aim, indication and medicines	Docetaxel + cyclophosphamide (TC) starts	Review response, toxicity and exposure	Complete planned course / next decision	Ongoing review / subsequent plan
Surgery EARLY	Adjuvant: surgery already completed	No automatic operation during cycles	Review postoperative recovery and symptoms	Surgery was completed before this adjuvant regimen	Pathology and recovery guide next steps
Radiotherapy EARLY	Assess indication	Usually after cytotoxic chemotherapy	Arrange sequential timing	Radiation team confirms start	Review effects / follow-up
Blood tests NCCP 00250	full blood count (FBC), kidney and liver profile	Repeat as clinically indicated; confirm the unit's pre-treatment test plan	Review testing needs at treatment changes	Follow active medicines and abnormalities	Follow unresolved abnormalities
Echo / heart NCCP 00250 · ESC	History / symptoms review	No routine TC-specific echo	Investigate if clinically indicated	Follow any cardiac concern	Individual plan
electrocardiogram (ECG) NCCP 00250	ECG only for clinical indications or interacting medicines; not a routine TC-cycle requirement.	Symptoms / risk-based repeat	Review if cardiac concern	ECG and echo answer different questions	Clinical indication
Cancer imaging NCCP 00250	Record baseline extent as appropriate	Clinical response / symptoms review	Review new symptoms; this TC regimen is adjuvant	Arrange radiotherapy or further treatment if indicated	Planned breast imaging; no routine computed tomography (CT)/positron emission tomography (PET)

Care / check	Before starting	Early cycles	Later cycles / transition	Next treatment / completion	Follow-up
Markers / bone health NCCP 00250	Tests selected for active medicines	No routine tumour-marker clearance	No routine blood tumour markers when well	 dual-energy X-ray absorptiometry (bone density) (DEXA) if endocrine/ bone-risk indication	Follow the endocrine and bone-health plan
Endocrine / bone BONE · EARLY	Confirm hormone receptors	Plan endocrine therapy when appropriate	Separate from this chemotherapy prescription	 Start/continue per multidisciplinary team (MDT); DEXA when indicated	Risk-based bone review

What each check is for

1 Before first dose

Baseline

Baseline blood count, kidney and liver tests. The team arranges repeat testing as clinically indicated and confirms which results are needed before treatment.

Ask / act: Confirm tests and consent before prescribing.

2 Heart-monitoring plan

Baseline → cycles → after treatment

TC contains no anthracycline or HER2-targeted antibody. There is no routine three-month echo programme for TC alone; arrange cardiac tests for symptoms, prior exposure or other clinical indications.

Ask / act: Name the team responsible for booking and reviewing results.

3

Treatment-phase review

Before each transition

Docetaxel and cyclophosphamide on Day 1 every 21 days for four to six cycles. The team confirms the planned course before treatment. Radiotherapy, when indicated, follows chemotherapy.

Ask / **act:** Confirm active medicines, support script and the next test checkpoint.

4

Response and follow-up

Planned review points

For early breast cancer, breast imaging and clinical review follow the agreed pathway. Routine CT/PET scans and blood tumour markers are not used to look for distant recurrence when there are no symptoms or clinical concerns.

Ask / **act:** Review symptoms and imaging together; do not change treatment on tumour markers alone.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Your practical treatment plan

STEP 1

Before the first visit

Blood tests, relevant cardiac assessment, medicines review, fertility discussion and the consent conversation come first.

STEP 2

Plan venous access

A cannula, peripherally inserted central catheter (PICC) or implanted port may be used. If a port is chosen, a trained team inserts it in a separate procedure with local anaesthetic and sometimes sedation; the team explains fasting, blood-thinner and wound instructions.

STEP 3

Infusion day

Staff check the line, give prescribed premedicines and treatment, and observe for reactions. Tell them immediately about burning, pain, rash or breathlessness.

STEP 4

Home and next review

This regimen does not routinely need a take-home infusion pump. Staff disconnect the infusion; a port is de-accessed when appropriate. Keep line-care and next-visit instructions.

Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

Visit time and transport

First visits can take longer because of loading doses, premedicines and observation. Ask the unit about expected time, transport and any escort needed after sedation or line insertion.

Phase-specific visit pattern

Infusion and observation times differ by drug and first versus later doses; the unit confirms the full visit duration, including blood review and preparation.

Checks across treatment

Your team arranges, reviews and acts on tests; an overdue result needs follow-up.

Blood tests

Baseline blood count, kidney and liver tests. The team arranges repeat testing as clinically indicated and confirms which results are needed before treatment.

Heart monitoring

TC contains no anthracycline or HER2-targeted antibody. There is no routine three-month echo programme for TC alone; arrange cardiac tests for symptoms, prior exposure or other clinical indications.

ECG

ECG only for clinical indications or interacting medicines; not a routine TC-cycle requirement.

Cancer scans

For early breast cancer, breast imaging and clinical review follow the agreed pathway. Routine CT/PET scans and blood tumour markers are not used to look for distant recurrence when there are no symptoms or clinical concerns.

Surgery, radiotherapy and later treatment

Surgery → this adjuvant chemotherapy → radiotherapy if indicated. The surgical team decides lumpectomy versus mastectomy; the radiation plan depends on surgery, pathology and individual risk.

Hormone-receptor-positive disease

Endocrine treatment is considered when appropriate for the cancer biology and setting. It has its own schedule and monitoring; it is not automatically part of this chemotherapy prescription.

Bone health

DEXA is linked to an aromatase inhibitor, ovarian suppression, treatment-induced menopause or other bone risk—not automatically to every chemotherapy course. The endocrine team confirms baseline testing and risk-based repeat scans.

Markers and follow-up

After local treatment, the team plans mammography of remaining breast tissue. A mastectomy side does not need routine mammography. Report new symptoms rather than waiting for a scan.

Genomic tests

Where appropriate, the MDT may discuss a validated genomic test to inform the chemotherapy decision. This regimen card does not calculate an Oncotype result or select treatment from a score.

SACTCHECK · CHAPTER 03

Symptoms and supportive care

Docetaxel + cyclophosphamide (TC) · NCCP 00250

TC

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

TC

Treatment or check	Timing in this view
Chemotherapy · Day 1	 **Day 1** intravenous (IV)
Administration reaction	 **Days 1–21** Report during or after any visit
Nausea / appetite	 **Days 1–5** May occur early or later
Blood counts / infection	 **Days 1–21** Risk varies; fever needs prompt advice
Fatigue / mouth changes	 **Days 1–21** Variable timing and severity
Nerves / skin / nails	 **Days 1–21** Can build across doses

No dates to enter: The chart explains the pattern. Your team books actual appointments and reviews results.

Treatment delays: A delayed dose does not automatically delay an echo or other test that is already due.

Phase changes: Do not change the schedule using this educational map.

> **Why these bands stay broad:** These broad symptom bands are educational possibilities, not measured risk curves. Infusion days are the planned schedule; delays require a new clinical plan. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00250 v9 · official regimen ↗

> **There are no guaranteed "safe days".** Fever, shivers, sudden illness, severe breathing changes or chest pain need prompt assessment on any day.

Between-treatment notes

Day 1–21: Your current reference cycle

Docetaxel and cyclophosphamide on Day 1 every 21 days for four to six cycles. The team confirms the planned course before treatment. Radiotherapy, when indicated, follows chemotherapy. Follow the team's appointment and medicine instructions; contact them about missed doses or new symptoms.

Support around the medicines

1  **The correct phase prescription**

TC uses moderate-risk antiemetic support.

2 **Premedicines and growth factors**

Docetaxel dexamethasone is taken for three days starting the day before treatment. Ask the team to reconcile this with antiemetic steroids.

3 **Tests and appointments**

The team should name the booking owner, due window and person reviewing each required test. Tell them if an expected appointment has not arrived.

4 **After chemotherapy**

Discuss recovery, fertility, rehabilitation and the next treatment or surveillance step.

Questions worth asking

When are heart checks due?

No routine fixed-cycle echo or ECG is mandated for this regimen alone; assess symptoms and individual cardiac risk. Test timing follows the treatment, clinical findings and local pathway.

Where do surgery and radiotherapy fit?

In adjuvant care, radiotherapy usually follows chemotherapy when indicated; endocrine therapy follows its own receptor-specific plan.

Do I need routine CT scans or tumour markers?

Early breast follow-up uses planned breast imaging. Routine CT / PET, blood tumour markers or other distant-recurrence screening are not recommended without symptoms or a clinical indication. After local treatment: annual mammography of remaining breast tissue; additional imaging when indicated.

Does a port mean a pump at home?

No. This regimen has no routine take-home infusion pump. A port is an access device; the team removes the needle after the visit when appropriate and arranges device removal separately.

Who arranges the scans and tests?

Your treating service should book the planned checks, review the results and explain any change. If an expected appointment is missing, contact the team; this map does not book tests or confirm that they have happened.

> **Try explaining the plan back:** Can you explain which medicines you are receiving now? What symptoms mean you should call urgently? Which blood tests, scans or heart checks come next, and who arranges them?

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Taxane / combination phase

Active medicines: Docetaxel + Cyclophosphamide

Emetogenic risk: Moderate · cyclophosphamide with docetaxel

Medicine	Dose / route	When
Ondansetron	8 mg orally twice daily	Day 1 NCCP alternative to 16 mg once daily; review QT / hepatic risks
Dexamethasone	8 mg orally	Day 1; reconcile docetaxel premedication Do not add to the docetaxel steroid course automatically
Dexamethasone	8 mg orally daily	Days 2–3 reference antiemetic row Reconcile the delayed period with docetaxel steroids; one final individualised steroid prescription
Metoclopramide	10 mg orally up to three times daily as required	Subsequent days: duration to be confirmed Review contraindications and renal function

Additional phase-specific support

- Docetaxel: dexamethasone 8 mg orally twice daily for 3 days starting the day before chemotherapy; at least 3 doses before the infusion. If missed, contact the team; any intravenous (IV) replacement is a consultant decision.
- Reconcile the total dexamethasone schedule with antiemetics, including aprepitant interactions. The listed antiemetic steroid is a reference row, not an instruction to add a second course.
- Assess granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) according to the exact regimen and individual risk.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hands / feet

Docetaxel

Report numbness or tingling before the next dose.

2 Blood / infection


Chemotherapy phase

Fever, shivers or suddenly unwell: call immediately.

3 Hair / scalp

Chemotherapy phase

Hair loss is common. Ask about scalp cooling before treatment where suitable.



4 Fluid / skin / nails

Docetaxel

Report swelling, painful skin or nail changes.

5 Mouth

Chemotherapy phase

Painful ulcers or difficulty drinking need advice.

6 Bladder

Cyclophosphamide

Report blood in urine or painful urination.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Report new breathing symptoms, severe diarrhoea, inability to drink, painful swollen limbs, unusual bleeding, or worsening numbness. Tell staff immediately about infusion-site pain or an allergic reaction.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

SACTCHECK · CHAPTER 04

Sources and further information

Docetaxel + cyclophosphamide (TC) · National Cancer Control Programme (NCCP) 00250

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **National Cancer Control Programme (NCCP) 00250 v9 · official regimen**
   https://healthservice.hse.ie/documents/6353/250_DOCEtaxel-cycloPHOS_TC.pdf
2. **ESMO 2024 early breast cancer guideline · treatment and follow-up**
   https://www.annalsofoncology.org/article/S0923-7534(23)05104-9/fulltext
3. **ESC 2022 cardio-oncology guideline · cardiac surveillance and risk**
   https://academic.oup.com/eurheartj/article/43/41/4229/6673995
4. **NICE NG101 · bone health and follow-up**
   https://www.nice.org.uk/guidance/ng101/chapter/Recommendations
5. **ESMO metastatic breast cancer guideline · 2026**
   https://www.annalsofoncology.org/article/S0923-7534(26)00902-6/fulltext
6. **eviQ · implanted port patient information**
   https://www.eviq.org.au/p/507
7. **eviQ · peripherally inserted central catheter (PICC) patient information**
   https://www.eviq.org.au/p/543
8. **HSE · chemotherapy side effects and urgent advice**
   https://www2.hse.ie/conditions/chemotherapy/side-effects/
9. **NOGG · section 7, aromatase inhibitors and fracture risk**
   https://www.nogg.org.uk/full-guideline/section-7-strategies-management-osteoporosis-and-fracture-risk
10. **NCCP NCIS antiemetics v8 (2025), Table 1**
    https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf
11. **ASCO · biomarkers in metastatic breast cancer**
    https://ascopubs.org/doi/10.1200/JCO.2015.61.1459