Weekly carboplatin + paclitaxel with radiotherapy

Chemotherapy and radiation treat stage III non-small-cell lung cancer when cisplatin is unsuitable. Your team explains your treatment aim and options.

Stage III non-small-cell lung cancer · concurrent chemoradiation · NCCP 00309 v6a · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Stage III non-small-cell lung cancer (NSCLC) weekly7-week regimen. This is distinct from oesophageal CROSS(5 weeks), metastatic chemotherapy and any separately prescribed consolidation.

My overall treatment

Paclitaxel then carboplatin by drip once every 7 days for seven weeks, alongside separately scheduled radiotherapy. Test results or side effects may change treatment. Any treatment after this course needs a separate decision and prescription.

7 day view

Treatment timing in this window

Paclitaxel then carboplatin

Day 1

Chemotherapy	Radiotherapy	After treatment
Both drugs at each weekly visit, after blood tests, symptom review and premedicines.	The radiation team sets the dose and appointments. A chemotherapy delay does not automatically stop radiation.	Your team reviews recovery, persistent symptoms and response. Scan or marker timing is individual.

A typical 7-day cycle

Day 1	Day 2–3	Day 4–7
Weekly treatment day	Support at home	Before the next weekly dose

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Blood count, kidneys, liver and GFR** Required by regimen	Measured GFR preferred; otherwise verified estimate
Before / during	**Hearing / nerves** If clinically indicated	When clinically indicated
During	**Blood count, kidneys and liver** Required by regimen	Weekly; creatinine within 48 hours for calculated GFR
After / review	**Recovery and response** Team-agreed plan	Team-agreed timing; no universal interval

Between treatments

- Use the prescribed anti-sickness medicines; report numbness, swallowing pain, fatigue or muscle and joint symptoms before treatment.
- Fever, new breathlessness, uncontrolled vomiting or inability to drink needs urgent advice, including after the course ends.

Confirm your prescription, radiation appointments and blood tests. Bring your medicine list and daytime / overnight contact numbers.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Nerves


Paclitaxel

Report numbness, tingling, weakness or difficulty with everyday tasks before treatment.

2 Mouth / swallowing

Chemoradiation

Tell the team if swallowing becomes painful. Inability to drink needs urgent advice.



3 Lungs / allergy

Chemoradiation

New breathlessness or worsening cough needs urgent review, including after treatment ends.

4 Sickness / appetite

Chemotherapy

Use prescribed anti-sickness treatment; call for uncontrolled vomiting or dehydration.

5 Kidneys / dose checks

Carboplatin

Kidney tests help calculate the dose. Report reduced urine or dehydration.

6 Blood cells / muscles

Chemotherapy

Call for fever or bleeding. Report muscle/joint pain and any infusion-site swelling.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology immediately for fever, shaking chills, sudden illness, new breathlessness, uncontrolled vomiting, inability to swallow fluids, much less urine or unusual bleeding. Severe breathing difficulty, chest pain, collapse or heavy bleeding needs emergency care. Tell staff immediately about infusion pain, swelling or allergy symptoms.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Stage III non-small-cell lung cancer · concurrent chemoradiation · NCCP 00309 v6a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Carboplatin + paclitaxel / RT

Both drugs once every 7 days for 7 weeks; radiotherapy appointments follow a separate plan.

Phase: _____ Planned weekly doses: 7

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My treatment record

Team stamps after each completed treatment.

W1	W2	W3	W4
Date / initials	Date / initials	Date / initials	Date / initials

W5	W6	W7
Date / initials	Date / initials	Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.