Topotecan by infusion · small-cell lung cancer

Check your exact treatment This patient guide covers the small-cell lung cancer indication and the stated route only.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Days 1–5 infusion support

Active medicines: topotecan

Emetogenic risk: low

Medicine	Dose / route	When
Individually prescribed sickness medicine	Single agent selected under low-risk parenteral guidance	Before each treatment dose; reconcile multi-day support No duplicated preventive courses. Product interactions and patient factors apply.

Additional phase-specific support

- Follow a written prescription with start/stop days.
- granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) may be used after individual risk assessment; it is not automatically prescribed.
- Monitor infusion site and reactions.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00311 v4 · pNCCP 00311 supportive care; NCCP antiemetic guidance · NCCP antiemetic guidance

Sources checked 2026-11-02 · SACTCheck v0.84.29 · Regimen-level drafts; prescriber and oncology-pharmacy verification required