BASELINE → TREATMENT → FOLLOW-UP

Checks across treatment

These columns show order of care, not equal time intervals. Blood counts are arranged weekly before treatment; kidney/liver tests repeat before each cycle. Scans and extra tests follow the consultant plan. There is no fixed universal scan interval.

Check	Before starting	Day 1	Days 2–3	Recovery / next cycle	Course review
Treatment and oral doses	Confirm extensive small-cell lung cancer (SCLC) and cisplatin unsuitability	Carboplatin infusion before oral etoposide	Oral etoposide twice daily on both days	No further etoposide until next authorised cycle	Continue only while benefit/ tolerability support it
Blood counts	Baseline blood profile	Review pretreatment results	Extra review if illness or bleeding	full blood count (FBC) weekly prior to treatment, as arranged	Follow any persistent low counts
Kidneys, liver and albumin	Renal/liver profile; measured GFR preferred	Verify kidney-based carboplatin dose	Report poor intake or low urine	Renal/liver profile before each cycle	Reassess after illness or organ change
Hearing and nerves	Baseline symptoms; audiology if indicated	Report new changes	Report tinnitus, numbness or functional problems	Regular hearing/neurological review	Follow symptoms that persist
Prevention, adherence and safety	Pharmacy capsule teaching; contact number	Dedicated carboplatin antiemetics; capsule sequence	Record capsules; follow reconciled antiemetics	Support plan covers two days after last systemic anticancer therapy (SACT); report missed/vomited doses	Review medicines and unused capsule return
Disease assessment	Confirm treatment aim	Clinical review	Report significant new symptoms	Tests/scans according to Consultant plan	Agree continuation, change or stop

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. National Cancer Control Programme (NCCP) 00319 v5 · official regimen

THE WHOLE TREATMENT JOURNEY

Carboplatin + Oral Etoposide: treatment and checks

Extensive-disease small-cell lung cancer · carboplatin + oral etoposide

Carboplatin is planned into a vein on Day 1, before oral etoposide. Etoposide capsules are prescribed twice daily on Days 1, 2 and 3 only, on an empty stomach. The cycle repeats every 21 days while treatment is helping and side effects remain acceptable. Your team decides when to delay, change or stop; the source does not set a fixed number of cycles.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Day 1	Days 2–3	Recovery / next cycle	Course review
Treatment and oral doses NCCP 00319	Confirm extensive SCLC and cisplatin unsuitability	Carboplatin infusion before oral etoposide	Oral etoposide twice daily on both days	No further etoposide until next authorised cycle	Continue only while benefit/ tolerability support it
Blood counts NCCP 00319	Baseline blood profile	Review pretreatment results	Extra review if illness or bleeding	FBC weekly prior to treatment, as arranged	Follow any persistent low counts
Kidneys, liver and albumin NCCP 00319	Renal/liver profile; measured GFR preferred	Verify kidney-based carboplatin dose	Report poor intake or low urine	Renal/liver profile before each cycle	Reassess after illness or organ change
Hearing and nerves NCCP 00319	Baseline symptoms; audiology if indicated	Report new changes	Report tinnitus, numbness or functional problems	Regular hearing/neurological review	Follow symptoms that persist
Prevention, adherence and safety NCCP 00319	Pharmacy capsule teaching; contact number	Dedicated carboplatin antiemetics; capsule sequence	Record capsules; follow reconciled antiemetics	Support plan covers two days after last SACT; report missed/vomited doses	Review medicines and unused capsule return
Disease assessment NCCP 00319	Confirm treatment aim	Clinical review	Report significant new symptoms	Tests/scans according to Consultant plan	Agree continuation, change or stop

What each check is for

1 Before the first cycle

Before treatment authorisation

Confirm disease, organ function and safe home dosing.

Ask / **act:** Confirm consent, capsule teaching and prevention plan.

Sources: NCCP 00319

2 After Day 3

After the last prescribed capsule doses

Avoid unintended continuation or replacement.

Ask / **act:** Record doses and follow support plan; contact team about omissions/vomiting.

Sources: NCCP 00319 · LEAFLET · ORALCARE

3 Before each new cycle

After recovery and review

Counts, organ function and tolerability may change.

Ask / **act:** Start another capsule course only after authorisation.

Sources: NCCP 00319

4 Course review

According to Consultant plan

Treatment should continue only while appropriate.

Ask / **act:** Discuss benefit, toxicity and next plan; return unused capsules.

Sources: NCCP 00319 · LEAFLET

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.29