Weekly cetuximab + FOLFIRI

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Day 8 · weekly cetuximab only

Active medicines: Cetuximab

Emetogenic risk: Low

Medicine	Dose / route	When
Antihistamine	As prescribed	At least one hour before cetuximab Cetuximab premedication remains applicable.
Corticosteroid	As prescribed	At least one hour before cetuximab Reconcile low-risk antiemesis with this corticosteroid; do not duplicate steroids.
Metoclopramide	10 mg PO up to three times daily as required	Rescue only as prescribed No automatic delayed steroid course is carried over from Day 1 chemotherapy.

Additional phase-specific support

- Cetuximab only: no irinotecan, oxaliplatin, fluorouracil or atropine is scheduled on Day 8.
- Continue skin care, electrolyte monitoring, infusion precautions and the urgent contact plan.
- Low-risk antibody-only antiemesis is distinct from Day 1 moderate-risk chemotherapy prophylaxis.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00328 v7 · p6 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-11-05 · SACTCheck v0.83.8 · Regimen-level drafts; prescriber and oncology-pharmacy verification required