BASELINE → TREATMENT → FOLLOW-UP

My treatment and safety checks

Routine safety tests are separate from scans and tumour markers, which follow your individual plan. No universal imaging or tumour-marker interval is specified.

Check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Treatment	Confirm combination plan	Day 1 docetaxel; Days 2–21 nintedanib	Capsules twice daily	After≥4 cycles, review possible continuation	No automatic phase switch
Blood count	Baseline	Before each cycle	Around nadir / if unwell	After last combination as indicated	Follow abnormalities
Kidneys / liver / blood pressure (BP)	Baseline bloods and BP	Bloods each cycle	BP as indicated; extra tests if needed	Check tolerance before change	Liver monthly if oral continuation
Scans / other care	Individual disease plan	No fixed scan interval	Report symptoms early	Assess benefit	Agreed follow-up

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. National Cancer Control Programme (NCCP) 00372 v6 · official regimen · Vargatef current EMA product information

THE WHOLE TREATMENT JOURNEY

Nintedanib + docetaxel: treatment and checks

Adenocarcinoma non-small-cell lung cancer (NSCLC) · combination phase

Day 1: docetaxel drip, with no nintedanib. Days 2–21: nintedanib capsules twice daily. Repeat every 21 days for up to 6 cycles (about 18 weeks without delays); treatment may stop earlier. A later tablet-only phase needs a new team-confirmed plan.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Treatment NCCP 00372 · SMPC	Confirm combination plan	Day 1 docetaxel; Days 2–21 nintedanib	Capsules twice daily	After≥4 cycles, review possible continuation	No automatic phase switch

Care / check	Before treatment	Treatment / review	Between visits	Phase decision	After treatment
Blood count NCCP 00372 · SMPC	Baseline	Before each cycle	Around nadir / if unwell	After last combination as indicated	Follow abnormalities
Kidneys / liver / BP NCCP 00372 · SMPC	Baseline bloods and BP	Bloods each cycle	BP as indicated; extra tests if needed	Check tolerance before change	Liver monthly if oral continuation
Scans / other care NCCP 00372 · SMPC	Individual disease plan	No fixed scan interval	Report symptoms early	Assess benefit	Agreed follow-up

What each check is for

1 Prepare

Before treatment

To help control adenocarcinoma lung cancer after first-line chemotherapy; treatment benefit and side effects are reviewed together.

Ask / act: Confirm the aim, phase, prescription, tests and contacts.

Sources: NCCP 00372

2 Review

Before each cycle / at agreed reviews

Check benefit and tolerability

Ask / act: Report symptoms and confirm the next plan.

Sources: NCCP 00372

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-02 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.30