Cobimetinib + vemurafenib — advanced melanoma

National Cancer Control Programme (NCCP) 00373 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

Usual starting doses are cobimetinib 60 mg once daily on days 1–21 and vemurafenib 960 mg twice daily, about 12 hours apart, continuously. Cobimetinib can be taken with or without food. Avoid regularly taking both daily vemurafenib doses on an empty stomach. Swallow both medicines whole and follow the separate labels.

STEP 2

Missed, vomited or held treatment

For cobimetinib, take a missed dose only if at least 12 hours remain before the next dose. For vemurafenib the interval is at least four hours. Otherwise skip the missed dose; do not double up. If you vomit after either medicine, do not take an extra dose. A hold or reduction may apply to one medicine or both: obtain a written component-specific restart plan.

STEP 3

Appointments and supply

Before each visit, check four things: the blood-test booking, how you will get there, your medicine supply and any help you need. Ask who will review the results and tell you whether the next dose can go ahead.

STEP 4

Paper copies and contacts

Keep the handbook, short guide and treatment card in one place. Ask your team to fill in the real daytime and emergency numbers and print a replacement when the plan changes. A phone is optional.

Sources: NCCP 00373 v3: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.