procarbazine, lomustine and vincristine (PCV) chemotherapy · 42-day cycle

Three chemotherapy medicines to help control glioma; the team confirms the exact diagnosis, treatment goal and cycle plan.


NCCP 00379: adjuvant treatment of grade II glioma after radiotherapy, or palliative treatment of recurrent high-grade glioma. · NCCP 00379 v4 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

NCCP 00379: adjuvant treatment of grade II glioma after radiotherapy, or palliative treatment of recurrent high-grade glioma. Use the exact schedule and indication agreed with your team.

My overall treatment

42 days: lomustine once D1; procarbazine D8–21; vincristine intravenous (IV) D8 + D29.



Course reviewed individually

Doses, starts and stopping are confirmed by the team; a card stamp is a record, not approval.

A typical 42-day cycle

Day 1–7	Day 8–21	Day 22–29	Day 30–42
One lomustine dose on Day 1	Daily procarbazine and first vincristine	Procarbazine finished; Day 29 infusion	Recovery before the next cycle

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Bloods / lung function** Required by regimen	Counts, kidney/liver, glucose and baseline lung-function assessment.
Before each treatment	**Safety bloods / symptoms** Required by regimen	Counts, kidney/liver; review bowels, nerves and any dose held.
Between / after	**Weekly counts / lungs / magnetic resonance imaging (MRI)** Product information / individual plan	Counts weekly ≥6 weeks after lomustine (SmPC); lung checks as indicated. MRI on your plan.

Between treatments

- Lomustine is taken once, not daily. Procarbazine ends on Day 21; review before the Day 29 infusion.
- Delayed low counts can occur after capsule days. Report fever, bleeding, bowel or nerve symptoms early.

Keep the three medicine days separate. Have the anti-sickness, bowel, food/alcohol and interaction instructions ready.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Brain / energy


Glioma / PCV

Fatigue or neurological changes need review. New seizures, sudden weakness or speech difficulty need emergency assessment.

2 Mouth / eating

PCV

Mouth soreness or appetite loss can affect intake. Keep mouth care and ask for help with pain, ulcers or difficulty eating.



3 Blood / infection

Lomustine / procarbazine

Counts may fall weeks after lomustine and worsen over cycles. Fever, sudden illness, bruising or bleeding needs urgent advice.

4 Lungs

Lomustine

New cough or breathlessness needs assessment. Lung injury can develop with treatment; keep baseline and any repeat lung tests.

5 Liver / kidneys

PCV

Attend organ-function blood tests. Report yellow skin or eyes, dark urine, reduced urine or dehydration rather than waiting for the next cycle.

6 Bowel / nausea

PCV / vincristine

Use the nausea and preventive bowel plan. Severe constipation, abdominal pain, vomiting or inability to pass wind needs urgent advice.

7 Nerves / movement

Vincristine

Report tingling, numbness, weakness, jaw pain or difficulty walking before the next infusion. Symptoms can affect hands, feet or other areas.

8 Fertility / pregnancy

PCV

Discuss fertility preservation and contraception before starting. Ask for written precautions for patients and partners; do not breastfeed during treatment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Use your oncology unit’s written emergency plan. Fever, shivering or suddenly feeling unwell needs urgent advice even on days without chemotherapy. Call 112 or 999 for a new seizure, sudden weakness or speech difficulty, loss of consciousness, severe breathing difficulty or uncontrolled bleeding. If you have an established seizure rescue plan, follow it and its emergency instructions. Contact the team promptly for a worsening headache, repeated vomiting, new cough or breathlessness, unusual bruising, yellow skin/eyes or symptoms that prevent drinking. Do not restart held chemotherapy without instructions.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

NCCP 00379: adjuvant treatment of grade II glioma after radiotherapy, or palliative treatment of recurrent high-grade glioma. · NCCP 00379 v4 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

PCV · 42 days

42 days: lomustine once D1; procarbazine D8–21; vincristine IV D8 + D29.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Ongoing review record; no fixed total is implied. The team authorises each new cycle.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.