Palbociclib with endocrine therapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Palbociclib with endocrine therapy · supportive-care plan

Active medicines: Palbociclib + Your endocrine partner

Emetogenic risk: Oral minimal–low

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Palbociclib is oral minimal–low emetic risk. Premedication is not usually required; agree rescue treatment if symptoms occur.
- Attend early and pre-cycle blood tests. Dose holds and reductions are clinician decisions. Growth-factor injections are not a routine automatic prescription for every patient.
- Agree help for hot flushes, joints, vaginal or sexual symptoms, sleep and bone health. Fulvestrant injection discomfort and any ovarian-suppression injections have separate support plans.
- Check whether the pack contains capsules or tablets and follow the food instructions for that product. Check new medicines, supplements and grapefruit interactions.
- Report mouth ulcers, persistent bowel symptoms, worsening fatigue, fever or new breathing symptoms promptly.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00414 v5a · p4 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.21 · Regimen-level drafts; prescriber and oncology-pharmacy verification required