# Gemcitabine + carboplatin AUC2

National Cancer Control Programme (NCCP) 00430 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Your practical treatment plan

Carboplatin AUC2 is given first, followed by gemcitabine 1,000 mg/m², on BOTH Days 1 and 8 of each 21-day cycle. Treatment continues until progression or unacceptable toxicity.

STEP 1

### Before the first visit

Blood tests, medicines review and the consent discussion come first. Review previous treatment, organ function and the treatment-day blood-test plan.

STEP 2

### Plan venous access

A cannula, peripherally inserted central catheter (PICC) or implanted port may be used. If a port is chosen, a trained team inserts it in a separate procedure with local anaesthetic and sometimes sedation; the team explains fasting, blood-thinner and wound instructions.

STEP 3

### Infusion day

Staff check the line, give prescribed premedicines and treatment, and observe for reactions. Tell them immediately about burning, pain, rash or breathlessness.

STEP 4

### Home and next review

This regimen does not routinely need a take-home infusion pump. Staff disconnect the infusion; a port is de-accessed when appropriate. Keep line-care and next-visit instructions.

### Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

### PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

### Visit time and transport

Allow time for blood review, preparation, treatment and observation. Your unit confirms the full visit duration. Arrange transport or an escort where advised.

### Phase-specific visit pattern

Carboplatin AUC2 is given first, followed by gemcitabine 1,000 mg/m², on BOTH Days 1 and 8 of each 21-day cycle. Treatment continues until progression or unacceptable toxicity. No take-home infusion pump is part of this schedule. Port de-accessing after a visit is different from removal of the implanted port.

Sources: NCCP 00430 v6 · official regimen · eviQ · implanted port patient information · eviQ · PICC patient information. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.