Panitumumab + modified FOLFOX-6

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Panitumumab + modified FOLFOX-6

Active medicines: panitumumab + oxaliplatin + folinic acid + fluorouracil

Emetogenic risk: moderate

Medicine	Dose / route	When
Ondansetron	16 mg orally	Day 1 before chemotherapy Dose may be adjusted for individual/QT risk.
Dexamethasone	8 mg orally daily	Days 1–3 Reconcile all steroid prescriptions.
Metoclopramide	10 mg orally up to three times daily as prescribed	As required on subsequent days Confirm suitability, duration and contraindications.

Additional phase-specific support

- No routine panitumumab premedication unless previous hypersensitivity/local policy indicates.
- Pre-emptive EGFR skin care and electrolyte monitoring; monthly Mg/Ca/electrolytes for two months after last panitumumab.
- Pump, line and cold-sensitivity education; diarrhoea rescue plan.
- No ice chips during fluorouracil in this oxaliplatin regimen. Review neuropathy and cold-triggered throat symptoms; new unexplained cough or breathlessness requires urgent assessment.
- If treatment is delayed for cytopenia, follow the source-specific weekly blood-count checks and maximum delay; assess the antibody and chemotherapy components separately when indicated.
- Medication review: NCCP contraindicates concomitant brivudine, sorivudine and analogues with fluorouracil; review dihydropyrimidine dehydrogenase (a medicine-processing enzyme) (DPD)-affecting medicines, phenytoin toxicity/levels and warfarin international normalised ratio (blood-clotting test) (INR).
- EMA Vectibix also requires interruption for intolerable dermatologic reactions; a grade below 3 does not automatically clear treatment.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00447 v7 · p6 · NCCP NCIS Medical Oncology antiemetics V8, Table 1

Sources checked 2026-11-05 · SACTCheck v0.84.35 · Regimen-level drafts; prescriber and oncology-pharmacy verification required