SACTCHECK · TREATMENT PASSPORT

# Nivolumab + ipilimumab · colorectal after chemotherapy

## A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Cycles 1–4: nivolumab 3 mg/kg, then ipilimumab 1 mg/kg, on Day 1 every 21 days. The first nivolumab-only dose is due three weeks after the final combination dose, subject to clinical review. Nivolumab alone: 240 mg intravenous (IV) or 600 mg under the skin every two weeks, or 1200 mg under the skin every four weeks.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

If the team changes the interval between nivolumab-only doses, they will give you a new appointment plan: the first four-weekly dose is two weeks after the last two-weekly dose, or the first two-weekly dose is four weeks after the last four-weekly dose. Do not change the schedule yourself. This does not change which formulations are permitted for your colorectal treatment.

https://sactcheck.com/patient/00551/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

# Nivolumab + ipilimumab · colorectal after chemotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record combination cycles 1–4, then the chosen nivolumab-only dose, route and interval. A completed stamp does not approve the next treatment.

**Combination cycle 1 · nivolumab then ipilimumab** · copy this sheet for another cycle of the same phase.

### Nivolumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | □ Usual breathing | □ New cough or breathlessness: call promptly | □ Symptoms with usual activity: urgent review | □ Breathless at rest: urgent assessment | □ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

### Ipilimumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Nivolumab + ipilimumab · colorectal after chemotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record combination cycles 1–4, then the chosen nivolumab-only dose, route and interval. A completed stamp does not approve the next treatment.

**Combination cycle 2 · nivolumab then ipilimumab** · copy this sheet for another cycle of the same phase.

### Nivolumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | □ Usual breathing | □ New cough or breathlessness: call promptly | □ Symptoms with usual activity: urgent review | □ Breathless at rest: urgent assessment | □ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

### Ipilimumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Nivolumab + ipilimumab · colorectal after chemotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record combination cycles 1–4, then the chosen nivolumab-only dose, route and interval. A completed stamp does not approve the next treatment.

**Combination cycle 3 · nivolumab then ipilimumab** · copy this sheet for another cycle of the same phase.

### Nivolumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | □ Usual breathing | □ New cough or breathlessness: call promptly | □ Symptoms with usual activity: urgent review | □ Breathless at rest: urgent assessment | □ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

### Ipilimumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Nivolumab + ipilimumab · colorectal after chemotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record combination cycles 1–4, then the chosen nivolumab-only dose, route and interval. A completed stamp does not approve the next treatment.

**Combination cycle 4 · nivolumab then ipilimumab** · copy this sheet for another cycle of the same phase.

### Nivolumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | □ Usual breathing | □ New cough or breathlessness: call promptly | □ Symptoms with usual activity: urgent review | □ Breathless at rest: urgent assessment | □ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

### Ipilimumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Nivolumab + ipilimumab · colorectal after chemotherapy

Cycle number: _____ Start date: _____ Next review: _____

Record combination cycles 1–4, then the chosen nivolumab-only dose, route and interval. A completed stamp does not approve the next treatment.

**Maintenance · confirm nivolumab dose, route and interval** · copy this sheet for another cycle of the same phase.

### Nivolumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | □ Usual breathing | □ New cough or breathlessness: call promptly | □ Symptoms with usual activity: urgent review | □ Breathless at rest: urgent assessment | □ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | □ Usual pattern | □ Up to 3 extra stools above normal | □ 4–6 extra stools/day or at night: call | □ 7+ extra stools/day or dehydration: urgent review | □ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | □ None | □ Mild change: report | □ Pain or daily tasks affected: call | □ Spreading infection or blistering: urgent review | □ Severe illness / collapse: emergency help |
| **Fatigue** | □ Usual energy | □ Tiredness improves with rest | □ Rest does not help; usual tasks affected: call | □ Self-care difficult: urgent review | □ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.