National Cancer Control Programme (NCCP) 00586 · clinician draft

Capecitabine + rectal radiotherapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Concurrent capecitabine and pelvic radiotherapy

Active medicines: capecitabine + radiotherapy

Emetogenic risk: oral_minimal_low

Medicine	Dose / route	When
Rescue antiemetic	Individual prescription	As required for oral capecitabine Radiotherapy nausea risk depends on field/site and is reviewed separately by the radiation team.

Additional phase-specific support

- No automatic high/moderate intravenous (IV) antiemetic course for capecitabine.
- Written tablet-day and missed/held-dose instructions; do not double doses or extend to weekends without the team's plan.
- Diarrhoea/dehydration, hand-foot and pelvic skin care; nutrition and urgent-contact support.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00586 v3 · pNCCP00586 supportive care; classification V6 Table 5 · NCCP NCIS Medical Oncology antiemetics V8, Table 1

Sources checked 2026-10-25 · SACTCheck v0.84.21 · Regimen-level drafts; prescriber and oncology-pharmacy verification required