Talazoparib · breast monotherapy

Talazoparib is taken once daily continuously until disease progression or unacceptable toxicity. The usual starting dose is 1 mg; kidney function, interactions and side effects can require a different prescription.

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Relative milestones, not equal time intervals. Your team books tests and reviews results; this map does not confirm completion.

Check	Before starting	First treatment	During treatment	Change / completion	Follow-up
Blood / organ function	full blood count (FBC), renal, liver	Early toxicity review	FBC / renal / liver every 4 weeks	Counts and restart review	Investigate prolonged cytopenia
Symptoms / medicines	Baseline assessment	Early symptom review	Keep the individual check plan	Reconcile at treatment changes	Report continuing symptoms

THE WHOLE TREATMENT JOURNEY

Talazoparib · breast monotherapy: treatment and checks

Germline BRCA1/2-mutated, HER2-negative locally advanced or metastatic breast cancer

Talazoparib is taken once daily continuously until disease progression or unacceptable toxicity. The usual starting dose is 1 mg; kidney function, interactions and side effects can require a different prescription.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First treatment	During treatment	Change / completion	Follow-up
Treatment plan NCCP 00605	gBRCA / HER2– eligibility	Daily capsules start	Daily without planned breaks	Until progression / intolerance	Next treatment plan
Blood / organ function NCCP 00605	FBC, renal, liver	Early toxicity review	FBC / renal / liver every 4 weeks	Counts and restart review	Investigate prolonged cytopenia

Care / check	Before starting	First treatment	During treatment	Change / completion	Follow-up
Symptoms / medicines NCCP 00605	Baseline assessment	Early symptom review	Keep the individual check plan	Reconcile at treatment changes	Report continuing symptoms
Cancer review METASTATIC · MARKERS	Confirm setting	Agree review plan	Response / symptom review	Next treatment / completion	Agreed surveillance

What each check is for

1 Before treatment

Before the first dose

Sources: NCCP 00605

2 Other checks

At the specified review points

Sources: NCCP 00605

3 Support and practical plan

At each treatment or phase change

Sources: NCCP 00605

4 Response and follow-up

Agreed review points

Sources: METASTATIC · MARKERS

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.