Paclitaxel Days 1, 8 and 15 · relapsed or refractory small-cell lung cancer (SCLC)

National Cancer Control Programme (NCCP) 00621 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Paclitaxel is given on Days 1, 8 and 15 of each 28-day cycle. Day 22 is a planned week off, not a fourth infusion. Treatment continues while helping and tolerated; there is no fixed cycle limit.

STEP 1

Before the visit

Confirm the next appointment on Days 1, 8 or 15. There is no Day 22 dose. Blood results and symptom review are needed before every dose.

STEP 2

At the unit

Paclitaxel is given through a cannula or suitable existing intravenous (IV) line over about one hour. The appointment also includes checks, preparation, premedication and observation and is longer than the infusion itself. This regimen has no routine take-home pump.

STEP 3

At home

Keep the contact number, medicine list and symptom record available. Ask who will arrange your next blood test, scan and review. Sedating medicines may affect driving.

Planned course

Paclitaxel is given on Days 1, 8 and 15 of each 28-day cycle. Day 22 is a planned week off, not a fourth infusion. Treatment continues while helping and tolerated; there is no fixed cycle limit.

Sources: NCCP 00621 v3 · official regimen · HPRA conventional paclitaxel SmPC; 9 January 2026. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.