National Cancer Control Programme (NCCP) 00688 · clinician draft

Atezolizumab + nab-paclitaxel

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Atezolizumab alone · after chemotherapy stops

Active medicines: Atezolizumab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine chemotherapy antiemetic or nab-paclitaxel premedication.
- Keep the source thyroid/blood monitoring and immune-toxicity review; stopping chemotherapy does not end immune risk.
- Confirm continuation has been authorised and use the atezolizumab schedule in this source.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00688 v2a · p7 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.19 · Regimen-level drafts; prescriber and oncology-pharmacy verification required