National Cancer Control Programme (NCCP) 00688 · clinician draft

Atezolizumab + nab-paclitaxel

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Days 1 and 15 · atezolizumab + nab-paclitaxel

Active medicines: Atezolizumab + Nab-paclitaxel

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally once	Before each administered chemotherapy dose Review QT risk, constipation, liver function and any steroid already prescribed.
Metoclopramide	10 mg orally up to three times daily as needed	Rescue duration to be confirmed Review contraindications, renal function and interactions.

Additional phase-specific support

- No routine corticosteroid or antihistamine premedication is required by the source for these medicines. Prior infusion reaction needs an individual plan.
- Give atezolizumab first, then nab-paclitaxel. Do not substitute solvent-based paclitaxel premedication.
- Review immune symptoms, count results and neuropathy before administration.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00688 v2a · p7 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.19 · Regimen-level drafts; prescriber and oncology-pharmacy verification required