weekly carboplatin AUC 2 / paclitaxel → dose-dense AC

Phase 1: four 21-day cycles; paclitaxel on Days 1, 8 and 15, carboplatin on all three days. Phase 2: four AC cycles every 14 days. Surgery and pathology determine the next step. See the whole-treatment map for monitoring and sequencing.

Source reconciled; independent oncology, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hair / scalp


Paclitaxel

Hair loss is common; ask about support and scalp cooling where suitable.

2 Nerves / nails

Paclitaxel

Report numbness, tingling or painful nails before the next dose.



3 Blood / infection

Carboplatin and paclitaxel

Fever, shivers, unusual bleeding or sudden illness: call immediately.

4 Infusion reaction

Carboplatin or paclitaxel

Tell staff immediately about flushing, chest tightness or breathing difficulty.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

This printed map shows phase 1. See the next page for all medicines and the online phase-specific maps. Report persistent effects after a phase ends.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Carboplatin

- Low blood counts, infection and bleeding
- Nausea, vomiting and fatigue
- Allergic reaction, sometimes after previous exposure
- Kidney or electrolyte changes

Paclitaxel

- Low blood counts and infection
- Numbness / tingling, sometimes lasting
- Hair loss, muscle or joint pain
- Infusion reaction
- Mouth, bowel, skin and nail effects

Doxorubicin

- Low counts, infection and mouth ulcers
- Nausea, hair loss and fatigue
- Heart damage, sometimes delayed
- Severe tissue injury if the infusion leaks
- Rare later blood cancers

Cyclophosphamide

- Low counts and infection
- Nausea, vomiting and hair loss
- Bladder irritation / bleeding
- Fertility effects and menopause
- Rare later blood cancers

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Separate anti-sickness plans

The platinum/taxane phase and AC need separate prescriptions. Carboplatin/paclitaxel is moderate emetic risk on each weekly treatment day. AC uses its own high-risk plan.

Growth-factor support

granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) is prescribed for every dose-dense AC cycle. Ask for the exact injection days and instructions.

Checks before the next dose

full blood count (FBC), renal and liver tests weekly throughout chemotherapy, including review before each AC dose. Report new symptoms before the next dose.

After chemotherapy

Surgery and pathology review follow chemotherapy. Further systemic treatment, radiotherapy and follow-up depend on the findings and the multidisciplinary team (MDT) plan; no routine postoperative immunotherapy is specified by this regimen.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00734/

Sources and review status are available online. Follow your own prescription and hospital instructions.