Bevacizumab + weekly paclitaxel

Bevacizumab 10 mg/kg is given on Days 1 and 15, with paclitaxel 80 mg/m² on Days 1, 8 and 15 of each 28-day cycle. Continue according to benefit, toxicity and the treating team's plan.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Blood pressure / vision


Bevacizumab

Severe headache or visual change needs urgent review.

2 Urine / kidneys

Bevacizumab

Protein checks can detect problems before symptoms; report swelling.



3 Abdomen / wounds

Bevacizumab

Severe abdominal pain or poor wound healing needs urgent advice.

4 Bleeding / clots

Bevacizumab

Report bleeding; sudden chest, breathing or stroke symptoms are emergencies.

5 Blood / infection

Paclitaxel

Fever, shivers or sudden illness: contact oncology.

6 Hands / feet

Paclitaxel

Report tingling or numbness before the next dose.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects; this is not a complete list. Report new or persistent symptoms even after treatment stops.

Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Bevacizumab

- High blood pressure / urine protein
- Bleeding or blood clots
- Delayed wound healing
- Rare bowel perforation / fistula
- Rare serious neurological effects
- Infusion reaction

Paclitaxel

- Low blood counts and infection
- Numbness / tingling, sometimes lasting
- Hair loss, muscle or joint pain
- Infusion reaction
- Mouth, bowel, skin and nail effects

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Blood pressure and urine

Attend the required checks even if you feel well. Report severe headache, visual change or swelling.

Operations and dental work

Tell oncology about planned procedures; bevacizumab timing and wound healing need review.

Paclitaxel premedicines

The first dose and later doses have different plans. A previous reaction changes the assessment.

Nerves and bleeding

Report numbness, unusual bleeding or bruising before the next visit.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00738/

Sources and review status are available online. Follow your own prescription and hospital instructions.