National Cancer Control Programme (NCCP) 00743 · clinician draft

# Eribulin · Days 1 and 15 of 28

**Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Eribulin · Days 1 and 15 of 28 · each treatment day

**Active medicines:** Eribulin

**Emetogenic risk:** Low

| Medicine | Dose / route | When |
|---|---|---|
| Ondansetron | 8 mg orally once | Each treatment day only<br>Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed |
| Metoclopramide | 10 mg orally up to three times daily as needed | Subsequent days: duration to be confirmed<br>Assess contraindications, renal function and interactions |

### Additional phase-specific support

- No routine hypersensitivity premedication is specified. Low-risk antiemetic support is reviewed separately, including QT interactions.
- This is not the licensed Day 1 / Day 8 schedule. The reason for the alternative schedule, evidence limits and off-label consent are part of the treatment discussion.
- Review interactions, QT risk and electrolytes before selecting an antiemetic; individualise rather than automatically issuing this draft.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00743 v3 · p4 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.19 · Regimen-level drafts; prescriber and oncology-pharmacy verification required