Phesgo + docetaxel

Phesgo and docetaxel on Day 1 every 21 days. Docetaxel is planned for at least six cycles; it may stop after 6–8 cycles or toxicity. Phesgo may continue alone while beneficial and tolerated.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Heart / breathing

Phesgo

Keep echoes; report breathlessness, swelling or reduced exercise tolerance.

2 Diarrhoea

Especially pertuzumab

Report watery stools early, particularly with poor intake or dizziness.

3 Injection reaction


Phesgo

Tell staff immediately about rash, chest tightness or breathing difficulty.

4 Thigh injection site

Phesgo

Report persistent or spreading redness, swelling or pain.



5 Hands / feet

Docetaxel

Report numbness or tingling before the next dose.

6 Blood / infection

Chemotherapy

Fever, shivers or suddenly unwell: call immediately.

7 Hair / scalp

Chemotherapy

Hair loss is common; ask about scalp cooling where suitable.

8 Mouth / appetite

Chemotherapy

Painful ulcers or reduced drinking need advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

The body map follows the medicines active in this phase.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Docetaxel

- Low blood counts, infection and sepsis
- Hair loss, mouth soreness and fatigue
- Numbness / tingling; sometimes lasting
- Fluid retention and nail changes
- Allergic infusion reaction

Pertuzumab

- Diarrhoea and dehydration
- Injection / allergic reaction
- Heart pumping-function reduction
- Rash and fatigue

Trastuzumab

- Heart pumping-function reduction / heart failure
- Injection or allergic reaction
- Diarrhoea and fatigue
- Rare serious lung effects

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Antiemetic plan

Taxane treatment uses a low-risk antiemetic plan; Phesgo alone uses a separate minimal-risk plan.

Premedicines

Docetaxel: prescribed dexamethasone starts the day before each dose. Ask for one reconciled steroid schedule.

Appointments and results

Echo before treatment and every 12 weeks (around cycle 5 if on time), including maintenance. Bloods before each cycle. Ask who books and reviews each test; report a missing appointment.

Phase transition

After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00796/

Sources and review status are available online. Follow your own prescription and hospital instructions.