Cisplatin, lomustine and vincristine (CLV)

CLV · 42 days NCCP 00806: selected adults with high-risk medulloblastoma or another eligible primitive neuro-ectodermal tumour, including pinealoblastoma. Use the exact prescription and schedule agreed with your specialist.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Days 8 and 15 · vincristine only

Active medicines: Vincristine

Emetogenic risk: Minimal

Medicine	Dose / route	When
Constipation prophylaxis	Agent / dose / duration to be completed by the prescriber and oncology pharmacist	Continue / adjust the written bowel plan during vincristine treatment Incomplete review row; not a prescription. Review constipation, ileus symptoms, peripheral/autonomic neuropathy and drug interactions before dosing.

Additional phase-specific support

- NCCP recommends preventive laxatives with vincristine. Report constipation, new tingling, weakness, balance or hand-function changes before each visit. Severe abdominal pain, vomiting or inability to pass stool or wind needs urgent assessment.
- Have all medicines checked, including seizure drugs, steroids, antibiotics, antifungals, non-prescription products and supplements. The team will confirm the local antifungal prevention plan and choose medicines that avoid vincristine interactions. Other infection prevention follows the individual specialist plan.
- Do not automatically repeat cisplatin hydration or its high-risk antiemetic course on vincristine-only days. Prior symptoms and the local individual plan can require additional support.
- Draft for clinician/pharmacist completion, not an issued prescription. Confirm agents, doses, timing, duration, interactions and rescue instructions with the current written plan.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

Sources checked 2026-11-05 · SACTCheck v0.84.39 · Regimen-level drafts; prescriber and oncology-pharmacy verification required