BASELINE → TREATMENT → FOLLOW-UP

# Tests and decisions across the treatment course

These are relative stages, not equal time intervals. Additional tests depend on symptoms and the selected chemotherapy source.

| Check | Preparation | Cycle 1 | Cycle 2 | Cycle 3 | Surgical review | Follow-up |
|---|---|---|---|---|---|---|
| **Treatment** | Confirm selected backbone | Day 1 all three; Day 8 gemcitabine | Repeat if appropriate | Final planned cycle | No automatic systemic anticancer therapy (SACT) continuation | Separate plan if further treatment needed |
| **Blood and organ tests** | full blood count (FBC), kidney, liver, glucose, thyroid and viral screen | FBC, kidney, liver, glucose; Day 8 counts | Repeat before treatment | Repeat before treatment | Recovery and fitness checks | Follow unresolved abnormalities |
| **Immune monitoring** | Document baseline symptoms | Symptoms before every dose | Thyroid every 3–6 weeks | Cortisol if clinically indicated | Report delayed symptoms | Vigilance at least five months |
| **Support and decisions** | Follow the written anti-sickness plan; Day 8 gemcitabine has its own review and support instructions. Carboplatin dosing uses a verified kidney-function measurement; report dehydration or a change in health. | Confirm written prevention plan | Review tolerability | Plan end of chemotherapy support | multidisciplinary team (MDT) agrees imaging and timing | Agree contacts and surveillance |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. National Cancer Control Programme (NCCP) 00849 v2 · official regimen · NCCP 00310 v8 · selected chemotherapy

THE WHOLE TREATMENT JOURNEY

# Nivolumab, gemcitabine and carboplatin before surgery: treatment and checks

**Neoadjuvant non-small-cell lung cancer (NSCLC) · selected gemcitabine / carboplatin backbone**

Up to three 21-day cycles. Day 1: nivolumab, gemcitabine and carboplatin. Day 8: gemcitabine alone after blood tests and review. Then surgical MDT reassessment. The five listed backbones are alternatives; this guide covers only your selected combination.

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

**Treatment, tests and reviews across the course**

| Care / check | Preparation | Cycle 1 | Cycle 2 | Cycle 3 | Surgical review | Follow-up |
|---|---|---|---|---|---|---|
| **Treatment** NCCP 00849 · COMPANION | Confirm selected backbone | Day 1 all three; Day 8 gemcitabine | Repeat if appropriate | Final planned cycle | No automatic SACT continuation | Separate plan if further treatment needed |
| **Blood and organ tests** NCCP 00849 · COMPANION | FBC, kidney, liver, glucose, thyroid and viral screen | FBC, kidney, liver, glucose; Day 8 counts | Repeat before treatment | Repeat before treatment | Recovery and fitness checks | Follow unresolved abnormalities |
| **Immune monitoring** NCCP 00849 · COMPANION | Document baseline symptoms | Symptoms before every dose | Thyroid every 3–6 weeks | Cortisol if clinically indicated | Report delayed symptoms | Vigilance at least five months |
| **Support and decisions** NCCP 00849 · COMPANION | Follow the written anti-sickness plan; Day 8 gemcitabine has its own review and support instructions. Carboplatin dosing uses a verified kidney-function measurement; report dehydration or a change in health. | Confirm written prevention plan | Review tolerability | Plan end of chemotherapy support | MDT agrees imaging and timing | Agree contacts and surveillance |

# What each check is for

## 1 Preparation

Before first infusion

Confirm the exact indication and backbone

**Ask** / **act:** Complete pathology, molecular, organ-function and consent review.

Sources: NCCP 00849 · COMPANION

## 2 Each cycle

Before each planned dose

Treatment safety can change

**Ask** / **act:** Review toxicity, results and the active support plan.

Sources: NCCP 00849 · COMPANION

## 3 Surgery assessment

After at most three cycles

Coordinate recovery and local treatment

**Ask** / **act:** Agree surgery timing; no automatic postoperative nivolumab.

Sources: NCCP 00849 · COMPANION

## 4 Follow-up

During and after surgery

Immune effects may be delayed

**Ask** / **act:** Keep the immune alert card and report new symptoms promptly.

Sources: NCCP 00849 · COMPANION

## At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-29 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.27