Fruquintinib (Fruzaqla) monotherapy

National Cancer Control Programme (NCCP) 00890 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Prescription and calendar

Take the prescribed fruquintinib dose once daily on Days 1–21, then take no fruquintinib on Days 22–28. Take it at about the same time each day, with or without food. Repeat only when your team authorises the next cycle.

STEP 2

Taking capsules

The usual starting dose is 5 mg once daily, but follow your own label. Swallow capsules whole with water, with or without food; do not chew, dissolve or open them. Tell the team before starting if swallowing capsules is difficult.

STEP 3

Missed or vomited dose

If a dose is less than 12 hours late, take it and then follow the usual schedule. If more than 12 hours late, skip it and take the next dose at the normal time. At exactly 12 hours, or if unsure, ask your pharmacist or team. Do not take a replacement capsule after vomiting; take the next dose at the usual time.

STEP 4

Tests and visits

Baseline and regular blood count, kidney/liver profile, thyroid function and urine protein. Blood pressure must be controlled before starting, then checked weekly for the first month and monthly thereafter, or more often if needed. For warfarin, international normalised ratio (blood-clotting test) (INR) weekly until stable, then before each cycle. Scans and markers follow the consultant plan.

STEP 5

Storage

Keep the original bottle tightly closed to protect from moisture and out of children’s sight and reach. No special temperature conditions are required. Follow the expiry date and ask the pharmacist how to return unused capsules; do not put them into household waste or wastewater.

STEP 6

Treatment changes

Your team must confirm dose changes and restarts. Bring the capsule pack, current calendar and medicine list to review.

Sources: NCCP 00890 v2a official fruquintinib regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.