Serplulimab, carboplatin and etoposide

National Cancer Control Programme (NCCP) 00908 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Appointments

Serplulimab is given first over about 60 minutes at the first visit. If tolerated, later doses may take about 30 minutes. Carboplatin takes about 30 minutes and intravenous (IV) etoposide about 60 minutes. Assessment, preparation and observation add time. Days 2–3 need further etoposide visits unless the separate oral alternative has been prescribed.

STEP 2

Route and phases

Induction lasts four planned 21-day cycles unless treatment needs to stop earlier. On Day 1 you receive serplulimab, carboplatin and etoposide by drip. With the standard IV schedule, etoposide is repeated on Days 2 and 3. Your team may instead prescribe etoposide by drip on Day 1 and capsules twice daily on Days 2 and 3. This is a separate alternative, not extra treatment. Oral and IV doses are not interchangeable milligram for milligram. Confirm which plan you have; take the capsules on an empty stomach as prescribed.

STEP 3

Maintenance visits

After induction, carboplatin and etoposide stop. Serplulimab alone can continue every three weeks while it is helping and its risks remain acceptable. There is no fixed universal number of maintenance cycles in this NCCP regimen.

STEP 4

Medicines and urgent advice

Keep the written anti-sickness schedule, capsule instructions if relevant, thermometer, immunotherapy alert card and oncology numbers available. Ask early about transport or help at home.

Sources: NCCP 00908 v1b · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.