Tucatinib + capecitabine + trastuzumab

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Tucatinib + capecitabine + trastuzumab

Active medicines: Tucatinib + Capecitabine + Trastuzumab

Emetogenic risk: Tucatinib / capecitabine oral minimal to low; trastuzumab minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Agree a personalised antiemetic and diarrhoea prescription; review CYP and other interactions. No repeated intravenous (IV) chemotherapy premedication is assumed for these tablets.
- Tucatinib continues every day; capecitabine is Days 1–14 then off Days 15–21. Confirm separate hold / restart instructions.
- Premedicines are not usually required. If prior hypersensitivity, consider paracetamol / antihistamine under the clinician plan.
- Explain when to hold capecitabine and call for diarrhoea, mouth soreness or painful hand-foot symptoms. Assess dehydration promptly.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00936 v1 · p10 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.21 · Regimen-level drafts; prescriber and oncology-pharmacy verification required