INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Ipilimumab alone — advanced melanoma

Understand your treatment. Know what to expect. Keep your plan to hand.

Displayed calendar
21 days
See the active treatment phase

Treatment stages
1
Explained in the treatment reference

Regimen source
NCCP 00105
Version 7

Reference treatment cycle

Treatment stages

01 **Treatment visits**
21-day reference cycle

Your team confirms the active stage and dates. The full treatment guide explains each step.

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	5
03	Symptoms and supportive care	9
04	Your record and contact card	15
05	Sources and further information	19

My name

Hospital / oncology team

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

Understanding your treatment

Ipilimumab alone — advanced melanoma · National Cancer Control Programme (NCCP) 00105

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.


What each medicine does

To help immune cells control advanced melanoma using ipilimumab alone; the planned infusion course is limited, but immune effects and follow-up continue beyond it.

CTLA-4 CHECKPOINT IMMUNOTHERAPY

Ipilimumab

Yervoy



Releases a CTLA-4 immune brake

Ipilimumab blocks CTLA-4, one of the signals that normally restrain T cells. This may help immune cells attack melanoma. The same immune activity can inflame healthy organs, sometimes after the last infusion; some hormone problems need lasting treatment.

Releases a CTLA-4 immune brake → Treatment plan and safety checks → Report new symptoms early

What to watch for

Report diarrhoea, blood in stools, severe headache, unusual weakness or spreading rash promptly.

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Understanding your diagnosis and treatment goal

Use your own pathology, stage and agreed plan alongside this guide.

Your diagnosis and stage

Record where the melanoma began, its pathology result and the sites involved now. This pathway treats melanoma that cannot be fully removed or has spread. Treatment after complete removal has a different aim and a separate plan.

The tumour test

The laboratory checks a sample of the melanoma for a BRAF V600 change. The result can identify a target for certain medicines. A change found in the tumour does not, by itself, establish an inherited condition.

Scans and local treatments

Add planned scans, any operation and any radiotherapy to the same calendar. Ask what each part is intended to achieve and which team arranges it.

New changes still matter

Tell your team about a changing mole, a persistent sore, a new lump or an unexplained symptom. Improvement in a known tumour does not explain a new problem elsewhere.

Your medicines and overall plan

The schedule

Ipilimumab by intravenous (IV) infusion once every 21 days, up to four doses. Complete within 16 weeks of the first dose, whichever limit comes first.

Important boundaries

This guide describes ipilimumab monotherapy at 3 mg/kg. It is not the nivolumab combination and does not automatically lead to nivolumab maintenance. A four-dose treatment course does not mean immune risks end after dose four.

Evidence and choices

Median overall survival for ipilimumab alone was 10.1 versus 6.4 months with gp100; HR 0.66, p=0.003. These are group results from that study, not a prediction for you. Historical gp100 comparator and selected previously treated population. Do not substitute the separate ipilimumab-plus-gp100 arm or compare indirectly with modern PD-1 treatment.

Decisions, support and useful records

Your priorities belong in the treatment discussion.

Benefits and alternatives

Ask what the treatment aims to achieve, what alternatives are suitable, how likely important benefits are in your situation and how treatment burden or uncertainty affects that choice. A trial average cannot promise an individual outcome.

Support alongside treatment

Supportive care addresses symptoms and daily life during cancer treatment. Palliative care can be part of that support; accepting it does not mean you must stop anticancer treatment.

A person you trust

Choose whether to involve a relative or supporter and what may be shared with them. Ask for an interpreter or a different information format. Check that you can explain the next dose, tests and urgent-contact plan in your own words.

Your document set

Keep the current treatment summary, medicine list, test plan and contact card together. Use the printed instructions when you need them; the QR code is an optional link to further information.

What the treatment evidence can and cannot tell you

Evidence needs to match the medicine, dose combination, disease stage and previous treatment.

SACTCHECK · CHAPTER 02

Your schedule, tests and appointments

Ipilimumab alone — advanced melanoma · NCCP 00105

Your treatment phases

Ipilimumab alone — advanced melanoma

21-day view



This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

21-DAY REFERENCE CYCLE

Treatment visits

Treatment visits **Repeat only as prescribed**

Ipilimumab by intravenous (IV) infusion once every 21 days, up to four doses. Complete within 16 weeks of the first dose, whichever limit comes first.

BASELINE → TREATMENT → FOLLOW-UP

Your testing and review plan

A reference cycle is not a personalised calendar. Tests listed as symptom-triggered or individual are not presented as mandatory tests at every visit. No universal imaging or tumour-marker interval is invented.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00105 v7: current regimen source

THE WHOLE TREATMENT JOURNEY

Ipilimumab alone — advanced melanoma: treatment and checks

Advanced melanoma

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Safety tests and symptoms NCCP 00105	Blood count, kidney and liver function, glucose and thyroid tests. Review autoimmune disease, previous immune reactions, transplantation, infection, pregnancy possibility and all medicines, including steroids.	Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, plus a careful symptom review. Cortisol and other tests are added when symptoms suggest endocrine or other immune effects. Normal routine blood results do not exclude colitis, lung, nerve or other inflammation.	Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.	Immune toxicity can start or persist after the four-dose course. Keep the alert card, follow-up appointments, steroid-taper or hormone-replacement plan and the individual scan schedule. Report new symptoms after treatment has ended.	Do not restart held treatment until the team confirms the medicine, dose and timing. Safety tests and unresolved symptoms are reviewed first.
Scans and tumour results NCCP 00105	Keep the pathology, stage and relevant biomarker report.	Response scans follow the individual disease plan, separate from safety bloods.	Report new symptoms rather than waiting for a scan.	Ask which scan, its planned date and who will explain the result.	A treatment hold or stop may change the next scan or follow-up date.
Written plan and contacts NCCP 00105	Record actual oncology and out-of-hours numbers; examples are not working contact details.	Bring your medicine list and record the dose actually given or taken.	Keep the wallet card and written urgent-symptom plan available.	Ask for a printed updated plan when treatment changes.	Keep the last-dose date and any steroid, hormone or other follow-up instructions.

What each check is for

1 **Treatment visits**

21-day reference cycle

To help immune cells control advanced melanoma using ipilimumab alone; the planned infusion course is limited, but immune effects and follow-up continue beyond it.

Ask / **act:** Confirm the current phase, prescription, tests and support medicines.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The source dose is ipilimumab 3 mg/kg over 90 minutes on day 1 of each 21-day cycle, with observation for one hour after the infusion. Allow additional time for symptom review, blood results and preparation. Report any previous infusion reaction before treatment.

STEP 2

Missed, vomited or held treatment

If a visit is missed or held, contact the unit. Do not count a cancelled visit as a dose received. The team reviews whether another dose remains appropriate within the four-dose and 16-week limits; do not assume a delayed dose can always be made up.

STEP 3

Appointments and supply

Before each visit, check four things: the blood-test booking, how you will get there, your medicine supply and any help you need. Ask who will review the results and tell you whether the next dose can go ahead.

STEP 4

Paper copies and contacts

Keep the handbook, short guide and treatment card in one place. Ask your team to fill in the real daytime and emergency numbers and print a replacement when the plan changes. A phone is optional.

Taking or receiving treatment safely

Keep the written prescription and actual treatment record together.

Other medicines

Give the team a complete list of prescribed, over-the-counter and herbal medicines, supplements and allergies. Ask before adding a new medicine; treatment interactions and symptom-masking can matter.

Practical arrangements

Write down where blood samples are taken, who receives the results and how you obtain the next appointment or medicine supply. Request a printed plan if that is easier to use.

After a treatment hold, change or completion

The next step should be written down, not assumed.

Follow-up and late effects

Immune toxicity can start or persist after the four-dose course. Keep the alert card, follow-up appointments, steroid-taper or hormone-replacement plan and the individual scan schedule. Report new symptoms after treatment has ended.

A treatment summary

Ask for a treatment summary showing what you received, important side effects, admissions, the last dose and the follow-up team. Share the current version with your GP and other clinicians.

Your next review

Before leaving, confirm the next test and review, how you will receive results and whom to call sooner. Keep those numbers available after treatment ends.

Symptoms and supportive care

Ipilimumab alone — advanced melanoma · NCCP 00105

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Treatment or check	Timing in this view
Treatment visit	 **Day 1** The source dose is ipilimumab 3 mg/kg over 90 minutes on day 1 of each 21-day cycle, with observation for one hour after the infusion. Allow additional time for symptom review, blood results and preparation. Report any previous infusion reaction before treatment.
Between-visit monitoring	 **Days 2–21** Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

Schedule and review: Ipilimumab by IV infusion once every 21 days, up to four doses. Complete within 16 weeks of the first dose, whichever limit comes first.

Holds, missed treatment and phase changes: If a visit is missed or held, contact the unit. Do not count a cancelled visit as a dose received. The team reviews whether another dose remains appropriate within the four-dose and 16-week limits; do not assume a delayed dose can always be made up.

Why these bands stay broad: This is a reference schedule. Your prescribed dose, blood results, symptoms and the team's written plan determine whether treatment goes ahead. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00105 v7: current regimen source ↗

There are no guaranteed "safe days". The prescribed dose, treatment decision and local urgent-contact plan take priority. Call 112 or 999 for severe breathing difficulty, severe chest pain, collapse, a new seizure, sudden weakness or speech difficulty, or uncontrolled bleeding. Follow your oncology unit's written urgent-contact plan for other new or worsening symptoms; do not wait for the next appointment.

Between-treatment notes

Day 1: Treatment day

The source dose is ipilimumab 3 mg/kg over 90 minutes on day 1 of each 21-day cycle, with observation for one hour after the infusion. Allow additional time for symptom review, blood results and preparation. Report any previous infusion reaction before treatment. Report symptoms and previous reactions before treatment.

Day 2–21: Between treatment visits

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy. Do not wait for the next appointment to report urgent symptoms.

Support around the medicines

1 Supportive medicines and symptom plan

NCCP classifies ipilimumab as low emetogenic risk; anti-sickness treatment is individual. Routine infusion premedication is not usually required, but previous reactions may change the plan. Do not start steroids or alter prescribed steroid tapers or hormone replacement without the team.

2 Skin and daily care

Use gentle skin care and sun protection. Report rash and itching early. Spreading rash, skin pain, blisters, peeling or sores affecting the mouth, eyes or genitals need urgent assessment. Continue melanoma skin self-checks and planned skin reviews.

3 Fertility, pregnancy and breastfeeding

Discuss pregnancy prevention, fertility and breastfeeding before treatment. Effective contraception and the duration after the last dose must be confirmed against the current ipilimumab product information and your written plan; this guide does not borrow the nivolumab five-month interval. Tell the team promptly if pregnancy is possible.

Tests, results and treatment decisions

Safety tests and scans answer different questions.

Before treatment

Blood count, kidney and liver function, glucose and thyroid tests. Review autoimmune disease, previous immune reactions, transplantation, infection, pregnancy possibility and all medicines, including steroids.

During treatment

Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, plus a careful symptom review. Cortisol and other tests are added when symptoms suggest endocrine or other immune effects. Normal routine blood results do not exclude colitis, lung, nerve or other inflammation.

What a result means

Ask which result is abnormal, whether the finding needs a repeat test and whether treatment is held, changed or stopped. Do not use a reference range or one normal result to decide independently that a dose is safe.

What to expect between visits

A quiet treatment day does not rule out later problems.



Supportive medicines

A useful daily record

Record symptoms, temperature when unwell, bowel changes, fluid intake when reduced, missed doses and any medicines used for relief. Note how symptoms affect eating, walking, sleep and normal activities; this helps the team judge severity.

When life is affected

Bring up difficulties with eating, sleep, pain, energy, work or caring for someone. The team can arrange symptom support while cancer treatment continues.

Skin, body changes and daily care

Use the detailed body map to locate symptoms, not to diagnose them.

The body map

The organ anchors show possible locations and warning symptoms. They do not show how common each effect is, and a symptom may have another cause. New symptoms should be assessed in their clinical context.

Food, fluids and activity

Choose food, fluids and movement that you can manage, within any personal restrictions. Ask for help early if you are losing weight, drinking less or finding everyday activity harder.

Keep checking the plan

A new medicine, fever, hospital admission, procedure or radiotherapy may change treatment timing. Tell the oncology team and take the wallet card or medicine list to other appointments.

Fertility, pregnancy and breastfeeding

Discuss these issues before the first dose whenever possible.

Fertility preservation

Ask whether treatment or the cancer may affect fertility and whether fertility preservation is appropriate before treatment. A change in periods or sexual function does not reliably establish infertility.

Written dates and methods

Record the agreed method, last treatment date and how long precautions continue. Different medicines have different intervals, so ask for clarification after a switch or combination change.

Sexual health

You can request a private discussion about intimacy, sexual symptoms or relationships. Ask for help if the agreed pregnancy precautions are difficult to follow.

Questions worth asking

Which tests are needed?

Blood count, kidney and liver function, glucose and thyroid tests. Review autoimmune disease, previous immune reactions, transplantation, infection, pregnancy possibility and all medicines, including steroids. Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, plus a careful symptom review. Cortisol and other tests are added when symptoms suggest endocrine or other immune effects. Normal routine blood results do not exclude colitis, lung, nerve or other inflammation. Immune toxicity can start or persist after the four-dose course. Keep the alert card, follow-up appointments, steroid-taper or hormone-replacement plan and the individual scan schedule. Report new symptoms after treatment has ended.

Can I use this without a phone?

Yes. The complete handbook, three-page summary and cut-out wallet card are printable. Ask the team to enter actual contact numbers and provide an updated paper plan when anything changes.

Try explaining the plan back: Can you explain your treatment setting, medicine schedule, next tests, the symptoms that need urgent advice and the number you would call?

The treatment this guide covers This guide describes ipilimumab monotherapy at 3 mg/kg. It is not the nivolumab combination and does not automatically lead to nivolumab maintenance. A four-dose treatment course does not mean immune risks end after dose four.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Ipilimumab alone — advanced melanoma · support plan

Active medicines: Ipilimumab

Emetogenic risk: Low per NCCP 00105 v7

Medicine	Dose / route	When
Ondansetron	8 mg orally once daily	Day 1, if prescribed under the low-risk plan Current NCIS V8 Table 1 p4. Review QT, electrolytes, constipation, hepatic dosing and interactions; this reference requires an individual prescription.
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days; duration to be prescribed Review contraindications, renal function, interactions and extrapyramidal adverse effects.

Additional phase-specific support

- The medicine list is a source-based low-risk reference for an individual consultant-approved/local prescription; it is separate from infusion-reaction premedication and immune-toxicity steroid treatment.
- NCCP classifies ipilimumab as low emetogenic risk; anti-sickness treatment is individual. Routine infusion premedication is not usually required, but previous reactions may change the plan. Do not start steroids or alter prescribed steroid tapers or hormone replacement without the team.
- Blood count, kidney/liver tests, glucose and thyroid tests before every cycle, plus a careful symptom review. Cortisol and other tests are added when symptoms suggest endocrine or other immune effects. Normal routine blood results do not exclude colitis, lung, nerve or other inflammation.
- Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.
- Discuss pregnancy prevention, fertility and breastfeeding before treatment. Effective contraception and the duration after the last dose must be confirmed against the current ipilimumab product information and your written plan; this guide does not borrow the nivolumab five-month interval. Tell the team promptly if pregnancy is possible.
- This guide describes ipilimumab monotherapy at 3 mg/kg. It is not the nivolumab combination and does not automatically lead to nivolumab maintenance. A four-dose treatment course does not mean immune risks end after dose four.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / hormones

Immunotherapy

Severe headache, dizziness, profound tiredness, thirst or altered vision can signal pituitary, adrenal, thyroid or glucose problems. Symptoms may need urgent hormone tests and treatment.

3 Heart

Immunotherapy

Rare heart inflammation can cause chest pain, palpitations, faintness or breathlessness. Severe chest pain, collapse or major breathing difficulty is an emergency.

5 Liver

Immunotherapy

Liver inflammation may first appear in blood tests. Yellow eyes or skin, dark urine, severe nausea or right-sided abdominal pain also need prompt review.

7 Nerves / muscles


Immunotherapy

New weakness, drooping eyelids, double vision, numbness or difficulty walking, swallowing or breathing may be a rare nerve or muscle effect. Seek urgent help; do not wait for the next visit.

9 Eyes

Immunotherapy

Eye pain, redness, light sensitivity or changed vision can indicate inflammation. Get prompt assessment; do not assume it is ordinary tiredness.



2 Lungs

Immunotherapy

New cough, shortness of breath or chest discomfort can mean lung inflammation or infection. Report changes promptly, even with normal recent blood results.

4 Skin

Immunotherapy

Rash and itching can occur. Rapid spreading, blistering, peeling or sores involving the mouth, eyes or genitals require urgent assessment.

6 Kidneys / urine

Immunotherapy

Kidney inflammation may be detected in blood tests. Report reduced urine, blood in urine, swelling or a marked change in thirst.

8 Bowels

Immunotherapy

New diarrhoea, blood or mucus, severe cramps or abdominal pain can mean bowel inflammation. Record changes from your usual pattern and call early; severe pain or bleeding needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Immunotherapy can inflame healthy organs during treatment or after it stops. Contact the oncology team promptly for new or worsening diarrhoea, blood or mucus in stools, abdominal pain, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell every treating clinician that you are receiving or have received immunotherapy.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

SACTCHECK · CHAPTER 04

Your record and contact card

Ipilimumab alone — advanced melanoma · National Cancer Control Programme (NCCP) 00105

Record each cycle separately. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Use a new record sheet for each cycle or review. Copy or print another sheet when needed; the number of sheets does not prescribe the length of treatment.

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

My diagnosis, plan and medicines

Complete these pages with your team. Leave unknown or inapplicable items blank. This record supports the consent discussion and your written prescription.

Diagnosis and overall plan

Original melanoma site / pathology date

Stage / Breslow thickness / ulceration

Lymph-node or other disease sites

BRAF result / date / other relevant tumour tests

Advanced or adjuvant intent / earlier surgery or radiotherapy (RT)

Medicines and safety

Anticancer medicines / actual dose / route / start date

Treatment received, missed or held / restart instruction

All other medicines / allergies / important previous reactions

Pregnancy-prevention or immune-alert plan / continuing precautions

Oncology / out-of-hours / pharmacy contacts

SACTCheck patient information. Your team confirms your personal plan. The treatment reference collects the clinical and medicine sources.

My checks, symptoms and support

Record what each test is checking and when the result will be discussed. Your team explains which examinations or scans are needed for your diagnosis and treatment.

Test and purpose	Date / booking contact	Result discussion	Agreed next step

Record bloods, skin/lesion examinations, scans and specialist checks separately. Tumour-marker and scan intervals are individually selected; do not infer them from the cycle calendar.

Symptoms, skin and support

New symptoms / onset / change / action taken

Lesion or wound / pain / eating / weight / daily function

Skin self-check plan / optional dated photograph record kept privately

Sun protection / practical, emotional or family support needs

Questions, response review and the outcomes that matter to me

My urgent contact plan / what my carer should do

SACTCheck patient information. Your team confirms your personal plan. The treatment reference collects the clinical and medicine sources.

Cycle number: _____ Start date: _____ Next review: _____

Record the dose actually taken or administered, symptoms, tests and agreed next step. A planned or held visit is not a completed treatment dose.

Ipilimumab alone — advanced melanoma · copy this sheet for another cycle of the same phase.

Ipilimumab · CTLA-4 checkpoint immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bowels	□ Usual pattern	□ New change: report	□ Persistent diarrhoea / constipation: call	□ Blood, severe pain or dehydration: urgent review	□ Collapse or severe sudden illness: 112/999
Abdominal pain	□ None	□ New pain: contact team	□ Persistent / worsening pain: urgent advice	□ Severe pain, blood or repeated vomiting: urgent assessment	□ Collapse or sudden severe illness: 112/999
Skin	□ No change	□ New small change: report	□ Persistent rash / itch: contact team	□ Painful, spreading or blistering rash: urgent review	□ Severe skin reaction with acute illness: emergency help
Headache	□ None	□ Mild and short-lived: record	□ New / persistent headache: call	□ Severe headache / visual change: urgent help	□ Sudden worst headache / weakness: 112/999
Weakness	□ Usual strength	□ New tiredness: record	□ New weakness limiting tasks: call	□ Marked weakness / faintness: urgent review	□ Sudden weakness, confusion or collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

Advanced melanoma · NCCP 00105 v7 · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Ipilimumab

Ipilimumab by IV infusion once every 21 days, up to four doses. Complete within 16 weeks of the first dose, whichever limit comes first.

Phase: _____ Planned cycles: 4

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1 Date / initials

C2 Date / initials

C3 Date / initials

C4 Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.42 · NCCP 00105 v7 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Sources and further information

Ipilimumab alone — advanced melanoma · National Cancer Control Programme (NCCP) 00105

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **National Cancer Control Programme (NCCP) 00105 v7: current regimen source**
   https://healthservice.hse.ie/documents/6724/105_V7_Ipilimumab_.pdf
2. **Hodi et al., NEJM 2010**
   https://pubmed.ncbi.nlm.nih.gov/20525992/
3. **NCI: fatigue during cancer treatment**
   https://www.cancer.gov/about-cancer/treatment/side-effects/fatigue
4. **NCI: melanoma — patient information**
   https://www.cancer.gov/types/skin/patient/melanoma-treatment-pdq
5. **NCI: melanoma treatment — clinical evidence**
   https://www.cancer.gov/types/skin/hp/melanoma-treatment-pdq
6. **NCI: eating and nutrition during cancer treatment**
   https://www.cancer.gov/about-cancer/treatment/side-effects/nutrition
7. **NCI: palliative care during cancer treatment**
   https://www.cancer.gov/about-cancer/advanced-cancer/care-choices/palliative-care-fact-sheet
8. **NCCP NCIS medical-oncology antiemetics V8, Table 1 p4, low risk**
   https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf