Pegylated liposomal doxorubicin · every 4 weeks

Pegylated liposomal doxorubicin on Day 1 every 28 days, typically for six cycles or until progression or unacceptable toxicity. The team confirms the intended duration and reviews benefit.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Palms / soles


Pegylated liposomal doxorubicin

Report painful redness, swelling or peeling early.

2 Mouth

Pegylated liposomal doxorubicin

Ulcers or difficulty eating / drinking need advice.



3 Heart

Pegylated liposomal doxorubicin

Keep electrocardiogram (ECG) / echo appointments; report breathlessness or swelling.

4 Blood / infection

Pegylated liposomal doxorubicin

Fever, shivers or sudden illness: call immediately.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects. The full discussion also includes fatigue, hair changes, fertility and other individual risks.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Pegylated liposomal doxorubicin

- Painful red palms / soles (hand-foot syndrome)
- Mouth soreness or ulcers affecting eating and drinking
- Low blood counts, infection, fatigue and nausea
- Heart damage and infusion reactions; risk depends partly on previous treatment

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Supportive medicines

No routine premedicines are specified. Low-risk antiemetic support is separate. Infusion reactions may require a different supervised plan for future doses.

Everyday care

Use skin care advised by your team; avoid heat, friction and tight shoes for 4–7 days after infusion. Report painful palms / soles or mouth ulcers early.

Tests and appointments

Ask who books each test, when it is due and who reviews the result. Tell the team if an expected appointment has not arrived.

Correct medicine and schedule

This is the pegylated liposomal formulation (for example Caelyx), not conventional doxorubicin or non-pegylated Myocet. These formulations and doses must not be interchanged.

Review the full medicine plan

Low emetic risk with an individual prevention/rescue plan; no routine hypersensitivity premedication is usually required. Reduce heat and friction affecting palms/soles; report painful redness, peeling or mouth ulcers before the next dose. Review cardiac symptoms and lifetime anthracycline exposure; a liposomal formulation still needs cardiac surveillance.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00205/

Sources and review status are available online. Follow your own prescription and hospital instructions.