Bevacizumab 7.5 · lung

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Bevacizumab component · combination or continuation

Active medicines: Bevacizumab

Emetogenic risk: Minimal; partner systemic anticancer therapy (SACT) separate

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine antiemetic prophylaxis for bevacizumab alone; select as-needed support under local policy.
- Do not use this component plan as the antiemetic plan for platinum chemotherapy.
- Premedication is not usually required unless a previous hypersensitivity warrants an individual plan.
- Blood-pressure treatment, urine-protein follow-up and procedure/wound timing need an individual plan.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00214 v8 · p5 · NCCP 00214 v 8 and current antiemetic classification

Sources checked 2026-11-02 · SACTCheck v0.84.30 · Regimen-level drafts; prescriber and oncology-pharmacy verification required