Eribulin · Days 1 and 8 of 21

Eribulin on Days 1 and 8 of each 21-day cycle. Day 15 is a planned treatment-free week. Treatment continues while beneficial and tolerated.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Hands / feet


Eribulin

Report numbness, tingling, weakness or loss of function.

2 Blood / infection

Eribulin

Fever, shivers or suddenly unwell: call immediately.



3 Heart rhythm

Eribulin

Fainting or new palpitations need urgent assessment.

4 Bowel

Eribulin

Report persistent constipation, abdominal pain or vomiting.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Selected medicine-specific effects. The full discussion also includes fatigue, hair changes, fertility and other individual risks.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Eribulin

- Low blood counts, infection and sepsis
- Numbness, tingling or weakness, sometimes lasting
- Fatigue, hair loss, nausea and constipation
- Changes to heart rhythm / QT in susceptible people

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Supportive medicines

No routine hypersensitivity premedication is specified. Low-risk antiemetic support is reviewed separately, including QT interactions.

Everyday care

Keep gently active as tolerated, pace rest and activity, and report nerve symptoms before the next dose.

Tests and appointments

Ask who books each test, when it is due and who reviews the result. Tell the team if an expected appointment has not arrived.

Correct medicine and schedule

EU dose labels use eribulin base. 1.23 mg/m² base corresponds to 1.4 mg/m² mesylate in some trial reports; these are not two alternative doses.

Review the full medicine plan

Low emetic risk; choose preventive/rescue medicines after considering QT prolongation and interactions. No routine hypersensitivity premedication is usually required. Review potassium/magnesium and correct abnormalities; monitor electrocardiogram (ECG) when cardiac/QT risks or symptoms warrant it. Severe neutropenia may be managed with granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) under the treating team. Avoid driving or machinery when tired or dizzy.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00228/

Sources and review status are available online. Follow your own prescription and hospital instructions.