Dabrafenib alone — advanced melanoma

National Cancer Control Programme (NCCP) 00237 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The usual starting dose is 150 mg twice daily, about 12 hours apart. Take capsules at least one hour before or at least two hours after food. Swallow whole with water. Use the prescribed strength and dose, and check the label again after any interruption or dose change.

STEP 2

Missed, vomited or held treatment

Skip a missed dose if the next dose is due in less than six hours. Do not take two doses together to make up for it. If you vomit after a dose, do not take a replacement dose; take the next dose when due. The team must give instructions for restarting after fever or another treatment hold.

STEP 3

Appointments and supply

Before each visit, check four things: the blood-test booking, how you will get there, your medicine supply and any help you need. Ask who will review the results and tell you whether the next dose can go ahead.

STEP 4

Paper copies and contacts

Keep the handbook, short guide and treatment card in one place. Ask your team to fill in the real daytime and emergency numbers and print a replacement when the plan changes. A phone is optional.

Sources: NCCP 00237 v6: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.