Adult advanced non-small-cell lung cancer (NSCLC) · ALK or ROS1 indication confirmed Adults with advanced ALK-positive or ROS1-positive NSCLC. National Cancer Control Programme (NCCP) 00243a is previously treated ALK disease; 00243b is ROS1 disease; 00243c is first-line ALK disease. Confirm the exact indication and local access; a listed indication, licence and reimbursement are different facts. This guide does not cover paediatric lymphoma or inflammatory myofibroblastic tumour.

NCCP 00243 · VERSION 7

Crizotinib monotherapy

The usual adult starting dose is 250 mg twice daily, every day, with no planned rest week. Use the strength and frequency on your own prescription. Treatment continues while it helps and side effects remain acceptable; there is no fixed course length.

Your treatment and review passport

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four sheets are reusable review periods, not a prescribed number of cycles. Follow the treatment and rest days agreed with your team; do not calculate a restart from this sheet.

These patient-language prompts are not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) assessment. Urgent symptoms override the form.

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Contact your oncology team immediately for new or worsening breathlessness or cough, fainting, palpitations, severe dizziness, rapidly increasing swelling/weight, sudden or severe loss of vision, yellow skin/eyes, dark urine, fever/shivering or sudden illness, severe abdominal pain, or persistent vomiting/diarrhoea with poor drinking or little urine. Do not wait for the next appointment. Severe breathing difficulty, collapse or severe chest pain needs emergency help.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

Your complete treatment reference

https://sactcheck.com/patient/00243/

Public information only. No personal information is encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Crizotinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Vision changes	☐	☐	☐	
Nausea	☐	☐	☐	
Diarrhoea	☐	☐	☐	
Constipation	☐	☐	☐	
Swelling	☐	☐	☐	
Fatigue	☐	☐	☐	
Breathlessness / cough	☐	☐	☐	
Palpitations / fainting	☐	☐	☐	
Skin / sun sensitivity	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Contact your oncology team immediately for new or worsening breathlessness or cough, fainting, palpitations, severe dizziness, rapidly increasing swelling/weight, sudden or severe loss of vision, yellow skin/eyes, dark urine, fever/shivering or sudden illness, severe abdominal pain, or persistent vomiting/diarrhoea with poor drinking or little urine. Do not wait for the next appointment. Severe breathing difficulty, collapse or severe chest pain needs emergency help. Emergency symptoms: 112 / 999.

Crizotinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Vision changes	☐	☐	☐	
Nausea	☐	☐	☐	
Diarrhoea	☐	☐	☐	
Constipation	☐	☐	☐	
Swelling	☐	☐	☐	
Fatigue	☐	☐	☐	
Breathlessness / cough	☐	☐	☐	
Palpitations / fainting	☐	☐	☐	
Skin / sun sensitivity	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Contact your oncology team immediately for new or worsening breathlessness or cough, fainting, palpitations, severe dizziness, rapidly increasing swelling/weight, sudden or severe loss of vision, yellow skin/eyes, dark urine, fever/shivering or sudden illness, severe abdominal pain, or persistent vomiting/diarrhoea with poor drinking or little urine. Do not wait for the next appointment. Severe breathing difficulty, collapse or severe chest pain needs emergency help. Emergency symptoms: 112 / 999.

Crizotinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Vision changes	☐	☐	☐	
Nausea	☐	☐	☐	
Diarrhoea	☐	☐	☐	
Constipation	☐	☐	☐	
Swelling	☐	☐	☐	
Fatigue	☐	☐	☐	
Breathlessness / cough	☐	☐	☐	
Palpitations / fainting	☐	☐	☐	
Skin / sun sensitivity	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Contact your oncology team immediately for new or worsening breathlessness or cough, fainting, palpitations, severe dizziness, rapidly increasing swelling/weight, sudden or severe loss of vision, yellow skin/eyes, dark urine, fever/shivering or sudden illness, severe abdominal pain, or persistent vomiting/diarrhoea with poor drinking or little urine. Do not wait for the next appointment. Severe breathing difficulty, collapse or severe chest pain needs emergency help. Emergency symptoms: 112 / 999.

Crizotinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Vision changes	☐	☐	☐	
Nausea	☐	☐	☐	
Diarrhoea	☐	☐	☐	
Constipation	☐	☐	☐	
Swelling	☐	☐	☐	
Fatigue	☐	☐	☐	
Breathlessness / cough	☐	☐	☐	
Palpitations / fainting	☐	☐	☐	
Skin / sun sensitivity	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Contact your oncology team immediately for new or worsening breathlessness or cough, fainting, palpitations, severe dizziness, rapidly increasing swelling/weight, sudden or severe loss of vision, yellow skin/eyes, dark urine, fever/shivering or sudden illness, severe abdominal pain, or persistent vomiting/diarrhoea with poor drinking or little urine. Do not wait for the next appointment. Severe breathing difficulty, collapse or severe chest pain needs emergency help. Emergency symptoms: 112 / 999.