Regorafenib (Stivarga) monotherapy

Targeted tablet treatment to help control previously treated colorectal cancer that has spread. Your team reviews benefit, side effects and your priorities.

Previously treated metastatic colorectal cancer · NCCP 00244 v5 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

This guide covers regorafenib for previously treated metastatic colorectal cancer. Regorafenib is also used in other cancers, but their eligibility and evidence are different. Confirm the treatment setting on your prescription.

My overall treatment

Days 1–21: prescribed tablets once daily after a light, low-fat meal. Days 22–28: no regorafenib. Repeat only when authorised by your team.

28 day view

Treatment timing in this window

Regorafenib tablets — Day 1–21

28-day cycle

Three tablet weeks, then one tablet-free week. No fixed total number of cycles.

Your treatment schedule

Days 1–21	Days 22–28
Once-daily prescribed tablets after a light, low-fat meal.	No regorafenib. Tests, recovery and next-cycle review.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline checks** Required / risk-based	Blood count, kidney, liver, thyroid and blood pressure. electrocardiogram (ECG)/QT if at risk.
Early	**Liver / blood pressure** Required by regimen	Liver: every two weeks for two months. blood pressure (BP): weekly for six weeks.
Later	**Ongoing checks** Required / individual	Liver monthly; BP before each cycle; blood count/kidneys regularly. Extra tests as indicated.
After / review	**Recovery / scans / markers** Individual plan	Team-agreed plan; follow unresolved effects. No fixed scan interval.

Between treatments

- Report painful hands/feet early; side effects can persist during the break.
- Same-day missed dose only; never two doses in one day. Do not replace a vomited dose.

Confirm the tablet calendar, meal and skin-care advice, first test dates and urgent contact numbers. If taking warfarin, arrange international normalised ratio (blood-clotting test) (INR) checks.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / energy

Regorafenib

Fatigue can affect daily life. Severe headache with confusion, visual change or a seizure needs emergency help.

3 Heart / blood pressure

Regorafenib

Blood pressure can rise without symptoms. Attend checks; chest pain, fainting or severe breathlessness needs emergency help.

5 Liver


Regorafenib

Liver injury may first appear in blood tests. Yellow skin or eyes, dark urine or confusion needs immediate advice.

7 Hands / feet / skin

Regorafenib

Pain, redness, blisters or thickened skin can limit walking or hand use. Report early and follow your skin-care plan.



2 Mouth / appetite

Regorafenib

Soreness, ulcers, taste changes or poor appetite can affect eating. Report difficulty drinking or weight loss.

4 Blood / infection

Regorafenib

Bleeding, bruising or infection can occur. Fever or sudden illness needs urgent advice; heavy bleeding needs emergency help.

6 Bowel / abdomen

Regorafenib

Diarrhoea can cause dehydration. Severe or persistent pain, black stools or blood in stools needs urgent help.

8 Kidneys / fluids

Regorafenib

Vomiting or diarrhoea can reduce fluid intake and urine. Very little urine or inability to drink needs urgent advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology immediately for fever, shivering, sudden illness, yellow eyes or skin, dark urine, severe or persistent abdominal pain, black or bloody stools, new bleeding, severe diarrhoea or inability to drink. Severe chest pain, major breathing difficulty, heavy bleeding, collapse, seizures, or a severe headache with confusion or visual change need emergency help: 112 / 999.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Previously treated metastatic colorectal cancer · NCCP 00244 v5 · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Regorafenib (Stivarga)

28 days · tablets Days 1–21 · no tablets Days 22–28 · team confirms each cycle.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.