# Trastuzumab subcutaneous (SC) · metastatic

National Cancer Control Programme (NCCP) 00272 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Your practical treatment plan

Trastuzumab 600 mg under the skin every 21 days, without a loading dose, while beneficial and tolerated. Any taxane or endocrine partner follows a separate prescription and monitoring plan.

STEP 1

### Before your visit

Confirm the dose, symptoms, required blood results and echo due window.

STEP 2

### Access and administration

SC trastuzumab needs no intravenous (IV) line for this medicine. Another medicine may still need a cannula or port.

STEP 3

### Treatment and observation

Injection over 2–5 minutes into alternate thighs. Observation is 30 minutes after the first injection and 15 minutes after subsequent injections, subject to approved local policy.

STEP 4

### After the visit

Contact the unit to rearrange a missed dose. SC doses remain at least 3 weeks apart; no loading dose. This medicine does not routinely require a take-home pump.

**Injection site**

Alternate thighs and avoid tender, bruised, red or hardened skin. Tell the nurse about a persistent or worsening local reaction.

**Visit length**

Injection over 2–5 minutes into alternate thighs. Observation is 30 minutes after the first injection and 15 minutes after subsequent injections, subject to approved local policy.

**Missed or delayed visit**

Contact the unit to rearrange a missed dose. SC doses remain at least 3 weeks apart; no loading dose.

Sources: NCCP 00272 v4 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.