INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Gemcitabine alone · non-small-cell lung cancer

Understand your treatment. Know what to expect. Keep your plan to hand.

Displayed calendar
28 days
See the active treatment phase

Treatment stages
2
Explained in the treatment reference

Regimen source
NCCP 00284
Version 5

Every 28 days

Treatment stages

01 **Repeated 28 day cycles**
3treatment visits per cycle

02 **Review benefit and recovery**
Next plan agreed

Your team confirms the active stage and dates. The full treatment guide explains each step.

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	3
03	Symptoms and supportive care	6
04	Sources and further information	10

My name

Hospital / oncology team

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

Understanding your treatment

Gemcitabine alone · non-small-cell lung cancer · National Cancer Control Programme (NCCP) 00284

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Call oncology immediately for fever or sudden illness, breathing difficulty, chest pain, uncontrolled vomiting, bleeding, very little urine, severe headache or confusion, or a spreading/blistering rash. Tell the nurse about infusion-site pain.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.


Your medicines

Gemcitabine aims to control lung cancer and its symptoms. Treatment is reviewed against side effects, your health and your priorities; benefit is not guaranteed.

CHEMOTHERAPY

Gemcitabine

Antimetabolite chemotherapy



Interferes with deoxyribonucleic acid (the cell's genetic material) (DNA) building

Gemcitabine interferes with making DNA, which dividing cancer cells need. Healthy blood-forming cells and other tissues can also be affected.

DNA building → Replication disrupted → Dividing cells affected

What to watch for

Report fever, unusual bleeding, breathlessness, rash, swelling or reduced urine promptly.

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Understanding the evidence

Gemcitabine has been studied in older adults with advanced lung cancer.

Small historical single-arm older-adult study cannot establish a comparative survival benefit or predict an individual outcome.

SACTCHECK · CHAPTER 02

Your schedule, tests and appointments

Gemcitabine alone · non-small-cell lung cancer · NCCP 00284

Your overall treatment

Treatment stages

01 **Repeated 28 day cycles**
3treatment visits per cycle

02 **Review benefit and recovery**
Next plan agreed

Your team confirms the active stage and dates. The full treatment guide explains each step.

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

3TREATMENT VISITS PER CYCLE

Repeated 28 day cycles

1 2 **Further cycles as agreed**

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

NEXT PLAN AGREED

Review benefit and recovery

Review

Your team decides when to scan and whether to continue, pause or change treatment. Surgery and radiotherapy are separate plans.

Lung treatment duration is individual; the pancreatic six-cycle cap is not transferred here.

BASELINE → TREATMENT → FOLLOW-UP

My tests and checks

Required safety checks are distinct from individual response imaging and follow-up. No routine tumour-marker interval is specified.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00284 v5 · official regimen

THE WHOLE TREATMENT JOURNEY

Gemcitabine alone · non-small-cell lung cancer: treatment and checks

Locally advanced or metastatic non-small-cell lung cancer (NSCLC) in older adults or people with ECOG2

Gemcitabine is given on Days 1, 8 and 15. Week 4 has no planned dose. Each cycle lasts 28 days. For lung cancer there is no fixed six-cycle limit in this source: continue only while the team considers treatment helpful and tolerable.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before treatment	Treatment days	Between doses	Next cycle	After treatment
Treatment NCCP 00284	Confirm lung indication	Gemcitabine Days 1/8/15	No dose Week 4	Repeat only if appropriate	Agree next plan
Blood tests NCCP 00284	full blood count (FBC), kidneys, liver	FBC before every dose	Extra tests if unwell	Renal/liver before cycle	Follow abnormalities
Other checks NCCP 00284	Audiometry/ creatinine clearance (CrCl) if indicated	Review symptoms	Call when unwell	Review benefit and toxicity	Individual scans/follow-up

What each check is for

1

Prepare

Before first dose

Gemcitabine aims to control lung cancer and its symptoms. Treatment is reviewed against side effects, your health and your priorities; benefit is not guaranteed.

Ask / act: Confirm aim, prescription, tests and contacts.

2

Review

Before each dose

Check treatment remains appropriate

Ask / act: Report symptoms and confirm blood results before dosing.

3 **Next plan**

At agreed review points

Assess benefit and recovery

Ask / **act:** Confirm scan and follow-up dates with the team.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before each visit

Bring your medicine list and report illness. The team reviews blood results before deciding the dose.

STEP 2

Three treatment visits

Day 1,Day 8 and Day 15 are planned infusion visits. Week 4 has no routine dose.

STEP 3

At home

Keep the prescribed support medicines and urgent contact plan available. Do not make up a missed dose yourself.

SACTCHECK · CHAPTER 03

Symptoms and supportive care

Gemcitabine alone · non-small-cell lung cancer · NCCP 00284

Day 1 → Day 28: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Treatment or check	Timing in this view
Gemcitabine only	 **Day 1** Day 1 only; after safety checks.
Gemcitabine only	 **Day 8** Day 8 only; after safety checks.
Gemcitabine only	 **Day 15** Day 15 only; after safety checks.
Symptom awareness	 **Days 1–28** Symptoms may occur on any day; this is not a personal risk curve.

Timing varies: Tiredness, bowel changes and infection can occur between visits. Blood tests and your symptoms guide dose decisions; there is no guaranteed safe day.

Changes to the plan: The team may delay, omit or reduce doses for safety. Do not add a missed dose yourself.

Why these bands stay broad: Relative treatment days, without personal dates. Symptoms are not confined to a fixed day. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00284 v5 · official regimen ↗

There are no guaranteed "safe days". Call oncology immediately for fever or sudden illness, breathing difficulty, chest pain, uncontrolled vomiting, bleeding, very little urine, severe headache or confusion, or a spreading/blistering rash. Tell the nurse about infusion-site pain.

Between-treatment notes

Day 1: Gemcitabine only

Treatment after blood-test and symptom review. Your team confirms whether this dose can proceed.

Day 2–7: At home

No planned chemotherapy on these days. Use only the support medicines prescribed for today. Call using your contact plan if unwell; do not wait for a treatment day.

Day 8: Gemcitabine only

Day 9–14: At home

Day 15: Gemcitabine only

Day 16–28: At home

Support around the medicines

1 **Sickness medicines**

Gemcitabine has low sickness risk. Follow the prescribed preventive or rescue medicine plan; do not take a platinum anti-sickness course for gemcitabine alone.

2 **Feeling unwell**

Do not dismiss fever as a flu-like drug effect. Call the oncology team immediately for fever, shaking chills or sudden illness.

3 **Medicines and fluids**

Follow your individual drinking advice. Report reduced urine, swelling, breathlessness or trouble drinking, and ask before new medicines or supplements.

Questions worth asking

Why are there three visits?

The same medicine is planned on Days 1, 8 and 15, followed by a week without a planned dose. Each visit has its own safety checks.

Is treatment always six cycles?

No. That fixed duration belongs to the separate pancreatic adjuvant indication. Your lung treatment continues only while appropriate.

When will I have scans?

Your consultant agrees timing from your disease and treatment goals; this regimen does not mandate one interval.

Try explaining the plan back: Explain back your treatment days, checks and whom to contact when unwell.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Gemcitabine alone · Days 1, 8, 15

Active medicines: gemcitabine

Emetogenic risk: low

Medicine	Dose / route	When
Dexamethasone	8 mg orally	Before each gemcitabine dose if prescribed NCIS low-risk reference; individualise and reconcile with any other steroid course.

Additional phase-specific support

- Gemcitabine has low sickness risk. Follow the prescribed preventive or rescue medicine plan; do not take a platinum anti-sickness course for gemcitabine alone.
- Do not dismiss fever as a flu-like drug effect. Call the oncology team immediately for fever, shaking chills or sudden illness.
- Follow your individual drinking advice. Report reduced urine, swelling, breathlessness or trouble drinking, and ask before new medicines or supplements.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / alertness


Gemcitabine

Urgent advice for severe headache, confusion, seizures or vision change.

2 Breathing

Gemcitabine

New or worsening breathlessness needs urgent advice.



3 Mouth / sickness

Gemcitabine

Report mouth sores, vomiting or trouble drinking.

4 Kidneys / fluids

Gemcitabine

Call for much less urine, swelling or sudden illness.

5 Skin

Gemcitabine

A spreading rash, blisters or peeling skin needs urgent review.

6 Blood cells

Gemcitabine

Fever, infection symptoms or unusual bleeding need prompt advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call oncology immediately for fever or sudden illness, breathing difficulty, chest pain, uncontrolled vomiting, bleeding, very little urine, severe headache or confusion, or a spreading/blistering rash. Tell the nurse about infusion-site pain.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

SACTCHECK · CHAPTER 04

Sources and further information

Gemcitabine alone · non-small-cell lung cancer · National Cancer Control Programme (NCCP) 00284

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **National Cancer Control Programme (NCCP) 00284 v5 · official regimen**
   https://healthservice.hse.ie/documents/6699/284_v5_Gemcitabine_Monotherapy_28_day.pdf
2. **Gemcitabine product information**
   https://www.medicines.org.uk/emc/product/3119/smpc
3. **Ricci et al. 2000: gemcitabine in older adults with advanced non-small-cell lung cancer (NSCLC)**
   https://pubmed.ncbi.nlm.nih.gov/10688490/
4. **NCCP Antiemetic Medicines for inclusion in NCIS (Medical Oncology) · v8, 10 February 2025**
   https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf