Osimertinib monotherapy

A targeted tablet for selected EGFR-positive lung cancers. Confirm whether your treatment follows surgery or treats advanced cancer.

EGFR-mutated non-small-cell lung cancer (NSCLC) · **adjuvant or advanced** · NCCP 00353 v5 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Take your prescribed dose once daily without a planned rest week. After surgery, treatment is planned for up to three years; advanced treatment continues while appropriate.

Your treatment pattern
Daily oral treatment without a rest week

Daily treatment
Keep taking the prescribed daily dose unless your oncology team changes it.

Reviews
Your team checks symptoms, medicines, blood results and benefit.

Continuous daily treatment

Every day	Review appointments
Take the dose on your prescription.	Tests and reviews do not create a tablet-free week.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**EGFR result; blood count, kidneys, liver and bone profile** Required by regimen	Confirm treatment setting and baseline blood tests
Before / early	**electrocardiogram (ECG) heart tracing** Required by regimen	Before starting and at the cycle-2 review; team confirms its date
During	**Blood count, kidneys, liver and bone profile** Required by regimen	Regular reviews; extra checks for symptoms or abnormal results
As indicated	**Further ECG / heart pumping assessment** If clinically indicated	Later ECGs and echo/multigated acquisition (heart pumping) scan (MUGA) when clinically indicated
After / review	**Recovery and cancer assessment** Team-agreed plan	Timing agreed with your team; report new symptoms promptly

Between treatments

- Do not create a tablet-free week around review appointments.
- Report new breathing difficulty, troublesome diarrhoea, eye symptoms or palpitations promptly.

Bring your medicine list and confirm the prescription, supply and missed-dose instructions.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Eye pain or blurred vision


osimertinib

Possible keratitis needs prompt specialist advice.

2 New cough or breathlessness

osimertinib

Possible lung inflammation needs urgent assessment.



3 Palpitations or fainting

osimertinib

Heart rhythm or pumping changes need assessment.

4 Diarrhoea

osimertinib

Report early; severe symptoms or dehydration need urgent advice.

5 Skin and nail changes

osimertinib

Blistering, peeling or mucosal sores are urgent.

6 Bruising or bleeding

osimertinib

Blood-count changes require tests and clinical review.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the oncology team urgently for new or worsening breathlessness, cough or fever; fainting, palpitations or chest pain; blistering/peeling skin or mouth/eye/genital sores; unusual bruising, bleeding or marked pallor; severe diarrhoea, dehydration or inability to drink. New eye pain, light sensitivity or blurred vision needs prompt advice. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

EGFR-mutated NSCLC · adjuvant or advanced · NCCP 00353 v5 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Osimertinib

Prescribed dose once daily, every day. Review dates do not create a treatment break.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.