INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Exemestane · daily hormone therapy

Understand your treatment. Know what to expect. Keep your plan to hand.

Treatment pattern
Ongoing
Follow the prescribed daily plan

Regimen source
NCCP 00376
Version 4

Continuous daily tablets

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	4
03	Symptoms and supportive care	8
04	Your record and contact card	13
05	Sources and further information	15

My name _____

Hospital / oncology team _____

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

Understanding your treatment

Exemestane · daily hormone therapy · National Cancer Control Programme (NCCP) 00376

How this medicine supports your treatment

After early breast cancer treatment, endocrine therapy reduces the risk of recurrence. In advanced disease, the aim is to control cancer and symptoms. Your setting determines the duration and follow-up.

HORMONE THERAPY · AROMATASE INHIBITOR

Exemestane

Reduces the oestrogen available to stimulate cancer cells

Aromatase normally makes oestrogen → The medicine blocks aromatase → Less oestrogen reaches cancer-cell receptors

Exemestane switches off aromatase, the enzyme that makes oestrogen in tissues after menopause. This reduces the hormone signal that can encourage hormone-sensitive breast cancer cells to grow. The body makes new enzyme over time, so tablets must still follow the prescribed daily schedule. It does not reliably switch off functioning ovaries; any ovarian suppression must be prescribed and checked separately.

Tell the team if joint pain, hot flushes, vaginal dryness or sleep disturbance affects daily life. Bone thinning may cause no early symptoms, so keep the agreed bone-health reviews.

The treatment being discussed

Who this information covers

Exemestane is used for selected postmenopausal hormone-sensitive breast cancer after surgery, as extended endocrine treatment or for advanced disease after earlier treatment. The team checks which previous endocrine medicine you received and why a switch or extension is appropriate. If switching after tamoxifen, the earlier years count towards the intended total treatment plan.

NCCP 00376 v4 · official regimen

Medicine and schedule

Take exemestane once daily continuously, preferably after a meal. There is no planned week off. Confirm the prior-treatment and duration plan. Swallow the tablet whole, preferably after a meal; do not chew or crush it. Skip a missed dose and continue at the next scheduled time; do not double the dose.

NCCP 00376 v4 · official regimen

Review and duration

Swallow the tablet whole, preferably after a meal; do not chew or crush it. Skip a missed dose and continue at the next scheduled time; do not double the dose. Count earlier endocrine treatment when reviewing an adjuvant switch. Extended treatment is a separate decision.

NCCP 00376 v4 · official regimen

Individual risks and support

The kidney and liver advice for exemestane differs from some other aromatase inhibitors: a routine dose reduction is not specified, but organ problems still need clinical review. Tell the pharmacist about all medicines and supplements because some can make exemestane less effective. Do not increase the dose yourself. Confirm menopause and any separate ovarian-suppression plan.

NCCP 00376 v4 · official regimen

Why this endocrine treatment is offered

For hormone-sensitive early breast cancer, endocrine treatment helps reduce the chance of the cancer returning. For advanced disease, it can help control cancer and symptoms.

Your likely benefit depends on your cancer, previous treatment and health. The treating team explains why this option fits your plan.

When to get help

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Report persistent limiting joint pain, new severe bone pain or a possible fracture, jaundice or marked new symptoms. Do not wait for the next scheduled review if you are becoming unwell.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

SACTCHECK · CHAPTER 02

Your schedule, tests and appointments

Exemestane · daily hormone therapy · NCCP 00376

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

Read the row for your treatment setting. Columns show order, not equal lengths of time. Agree the due window, booking team, result reviewer and whom to contact if a check is missing.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00376 v4 · official regimen · HPRA Exemestane SmPC cited by NCCP · Pan-Asian adapted ESMO early breast cancer guidelines · Pan-Asian adapted ESMO metastatic breast cancer guidelines · NICE NG 101: early and locally advanced breast cancer · ESO-ESMO ABC 5 consensus: tumour-marker interpretation · IOF/CABS/ECTS/IEG/ESCEO/IMS/SIOG updated joint statement on aromatase-inhibitor-associated bone loss

THE WHOLE TREATMENT JOURNEY

Exemestane · daily hormone therapy: treatment and checks

Your treatment pattern

Continuous daily tablets

This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

Postmenopausal ER-positive breast cancer after prior endocrine treatment

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First weeks	During treatment	Longer-term review	Change / completion
Treatment plan NCCP 00376	Confirm indication and prescription	Start the agreed daily / injection schedule	Continue unless prescriber changes the plan	Review duration, benefit and tolerability	Agree continuation, switch or completion
Bloods / symptom review NCCP 00376	full blood count (FBC), renal and liver profile	Review tolerability and medicines	Further tests when clinically indicated	Review symptoms, adherence and results	Follow unresolved abnormalities
electrocardiogram (ECG) / echo NCCP 00376	No routine ECG / echo for this medicine alone	Investigate new cardiac symptoms	Further cardiac tests by risk / symptoms	Companion-drug cardiac rules remain separate	Follow abnormalities when indicated

Care / check	Before starting	First weeks	During treatment	Longer-term review	Change / completion
Bone density / dual-energy X-ray absorptiometry (bone density) (DXA) NCCP 00376 · AIBL2025	Adjuvant aromatase inhibitor (AI): baseline DXA unless already on adjuvant bisphosphonates	Review previous DXA and fracture risk	No cycle-based DXA appointments	Lower-risk AI: reassess in 1–2 years; individualise	Continue bone care according to risk
Early-breast surveillance NCCP 00376	Confirm early / curative setting	Set first follow-up mammogram date	Assess new symptoms promptly	Annual mammography of remaining breast tissue	No routine computed tomography (CT) / markers if asymptomatic
Advanced-disease response NCCP 00376	Document disease sites / baseline assessment	New symptoms can bring assessment forward	Generally every 2–4 months; individualise	Adapt interval and imaging to disease / benefit	Markers supplement, never replace, assessment
Early: radiotherapy ESMO_EBC2024	Review surgery / pathology	Plan radiotherapy if indicated	Coordinate endocrine timing	Follow the early-breast plan	Review late effects when relevant

What each check is for

1 Confirm the plan

Before starting or changing treatment

Exemestane is used for selected postmenopausal hormone-sensitive breast cancer after surgery, as extended endocrine treatment or for advanced disease after earlier treatment. The team checks which previous endocrine medicine you received and why a switch or extension is appropriate. If switching after tamoxifen, the earlier years count towards the intended total treatment plan.

Ask / act: Confirm treatment setting, medicines, menopause and the planned review.

2 Checks and symptoms

At the agreed review points

FBC, renal and liver profile, confirmation of menopause when uncertain, and bone / fracture assessment. Consider baseline vitamin D as specified by the product information. The NICE adjuvant baseline-DXA recommendation has an exception for patients already receiving adjuvant bisphosphonates. Review symptoms, adherence and bone health. Repeat bone density and lipids when indicated; assess other blood tests according to the clinical plan.

Ask / act: Agree who books tests, reviews results and responds to new symptoms.

3

Review ongoing benefit

At setting-specific decision points

After early breast cancer treatment, endocrine therapy reduces the risk of recurrence. In advanced disease, the aim is to control cancer and symptoms. Your setting determines the duration and follow-up.

Ask / act: Review duration, alternatives and patient priorities; do not stop or extend treatment automatically.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Checks throughout treatment

Your team arranges tests and explains the results. A planned check is not confirmation that it has been completed.

Treatment-specific checks

FBC, renal and liver profile, confirmation of menopause when uncertain, and bone / fracture assessment. Consider baseline vitamin D as specified by the product information. The NICE adjuvant baseline-DXA recommendation has an exception for patients already receiving adjuvant bisphosphonates. Review symptoms, adherence and bone health. Repeat bone density and lipids when indicated; assess other blood tests according to the clinical plan.

NCCP 00376 v4 · official regimen

Heart checks

ECG or other heart tests depend on symptoms and personal risk; this medicine alone does not require a routine HER2-style echocardiogram schedule.

NCCP 00376 v4 · official regimen

Bone health

Offer baseline DXA when starting an adjuvant aromatase inhibitor if not already receiving adjuvant bisphosphonates; assess fracture risk and relevant previous results. Under the 2025 joint guidance, patients with a T-score above –2.0 and no additional risk factors generally have risk and bone density reassessed every one to two years. Patients on bone-protective treatment generally have BMD reassessed about every two years. Fracture, substantial bone loss or a change in risk can require earlier review. The clinician selects the interval and any bone treatment; these are not automatic yearly scans for everyone.

NICE NG 101: early and locally advanced breast cancer · IOF/CABS/ECTS/IEG/ESCEO/IMS/SIOG updated joint statement on aromatase-inhibitor-associated bone loss

Your setting changes the pathway

Choose the pathway that your team has confirmed; these are not all extra treatments you must have.

Early-breast follow-up

After curative treatment, annual mammography is generally used for remaining breast tissue: both breasts after conservation or the opposite breast after unilateral mastectomy. The mastectomy side is not routinely mammographed. Routine CT, bone scans and tumour markers are not recommended for asymptomatic early-breast surveillance. New symptoms need assessment between appointments.

Pan-Asian adapted ESMO early breast cancer guidelines · NICE NG 101: early and locally advanced breast cancer

Surgery, chemotherapy and radiotherapy

In an early-breast pathway, chemotherapy, if indicated, precedes endocrine treatment. Surgery and its pathology guide adjuvant choices. Postoperative radiotherapy is usually delivered after recovery and any adjuvant chemotherapy; aromatase inhibitors may overlap radiotherapy under the agreed plan. Tamoxifen timing around radiotherapy is agreed by the treating teams. None of these additional treatments is automatically required by an endocrine prescription.

Pan-Asian adapted ESMO early breast cancer guidelines · eviQ 3650 v 6: adjuvant whole-breast radiotherapy



Advanced-disease scans

For advanced breast cancer, response assessment is generally every two to four months, adapted to disease pace, sites, symptoms and treatment. Indolent disease may be reviewed less often; new symptoms can prompt earlier scans. The team chooses the imaging method; this is not a universal three-month CT chest/abdomen/pelvis order.

Tumour markers

Tumour markers may supplement imaging and clinical assessment in advanced disease if they are informative, especially if elevated at baseline. They do not replace scans, and a marker change alone should not determine treatment failure. There is no universal fixed marker timetable.

Menopause, pregnancy and contraception

Confirm the postmenopausal endocrine state. Any ovarian-suppression combination needs a separate prescription and monitoring plan.

SACTCHECK · CHAPTER 03

Symptoms and supportive care

Exemestane · daily hormone therapy · NCCP 00376

Your endocrine treatment and review milestones

These stages show the order of care, not equal lengths of time. No dates or personal details are needed. Follow the prescribed tablet schedule between reviews; the treatment setting determines when it is reviewed or completed.

Treatment or check	Timing in this view
Daily medicine	**First weeks → During treatment → Longer-term review → Change / completion** 25 mg orally once daily after a meal · continuous unless changed by prescriber
Before the first dose	**Before starting** Confirm menopause, treatment setting, baseline bloods, medicines and bone-health plan
Symptoms and tolerability	**First weeks → During treatment → Longer-term review → Change / completion** May persist or change; report troublesome symptoms early
Planned reviews	**During treatment → Longer-term review → Change / completion** Review symptoms, adherence, bone health and the planned duration or disease response

Swallow the tablet whole, preferably after a meal; do not chew or crush it. Skip a missed dose and continue at the next scheduled time; do not double the dose.

Count earlier endocrine treatment when reviewing an adjuvant switch. Extended treatment is a separate decision.

> Columns show sequence, not equal durations. These are reference milestones, not booked appointments. Symptom bands do not predict severity or a safe time to ignore symptoms.

Practical support

Symptoms and daily life

Support medicines

No routine antiemetic, steroid premedication or granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) is required by this endocrine monotherapy. Ask for individual symptom support.

Tests and appointments

Ask who books each check, when it is due and who reviews it. Tell the team if an expected appointment has not arrived.

Taking tablets / attending injections

Questions to ask

When are bloods, ECGs or echoes needed?

FBC, renal and liver profile, confirmation of menopause when uncertain, and bone / fracture assessment. Consider baseline vitamin D as specified by the product information. The NICE adjuvant baseline-DXA recommendation has an exception for patients already receiving adjuvant bisphosphonates. Review symptoms, adherence and bone health. Repeat bone density and lipids when indicated; assess other blood tests according to the clinical plan. There is no routine ECG or echo schedule for exemestane alone. Investigate symptoms and manage cardiovascular risk when indicated.

Which scans apply to me?

After curative treatment, annual mammography is generally used for remaining breast tissue: both breasts after conservation or the opposite breast after unilateral mastectomy. The mastectomy side is not routinely mammographed. Routine CT, bone scans and tumour markers are not recommended for asymptomatic early-breast surveillance. New symptoms need assessment between appointments. For advanced breast cancer, response assessment is generally every two to four months, adapted to disease pace, sites, symptoms and treatment. Indolent disease may be reviewed less often; new symptoms can prompt earlier scans. The team chooses the imaging method; this is not a universal three-month CT chest/abdomen/pelvis order.

How long will treatment continue?

If you switch after tamoxifen, the earlier years count towards the intended total endocrine course. Switching does not automatically mean starting a new full course. The team confirms your intended total and whether a later extension is appropriate. Advanced-disease treatment continues according to benefit and tolerance.

Do I need a port or home pump?

No port or home infusion pump is needed for this medicine itself. Any separately prescribed medicine has its own access requirements.

Can I use HRT, vaginal oestrogen or supplements?

Check with the oncology team before starting any hormone-containing treatment, including vaginal oestrogen, or any new medicine or supplement. Some can oppose this treatment or change how it works. The team can suggest symptom treatments and decide whether specialist review is needed.

What if periods return or menopause is uncertain?

Tell the team promptly about a return of periods or new vaginal bleeding. No periods after chemotherapy does not always mean the ovaries have permanently stopped working. Hormone blood tests may be needed. Ovarian suppression, if prescribed, has its own schedule and monitoring.

What should I do if symptoms make tablets difficult to continue?

Tell the team early about joint pain, hot flushes, sleep, vaginal or sexual symptoms, fatigue or missed tablets. Support, a supervised break or a change of medicine may be possible. Agree the plan with the prescriber rather than changing or stopping treatment yourself.

Can another medicine make exemestane less effective?

Yes. Some medicines for seizures or infections, and St John's wort, can reduce its effect. Check every new prescription, over-the-counter medicine and supplement with the oncology pharmacist. Do not compensate by taking extra exemestane.

Do I need calcium and vitamin D?

The NCCP regimen advises daily calcium and vitamin D supplementation. Ask the team or pharmacist to confirm the suitable product and dose, taking account of your diet, blood results, kidney health and any supplements you already take. Bone-protective medicines and bone-density checks follow your individual fracture-risk plan.

Explain the plan back: What is your treatment schedule, which checks apply to your setting, and who books and reviews them?

Take exemestane once daily continuously, preferably after a meal. There is no planned week off. Confirm the prior-treatment and duration plan.

> Clinician supportive-care discussion draft. This is not an issued prescription.

Minimal emetogenic risk

No routine prophylactic antiemetic, corticosteroid premedication or granulocyte colony-stimulating factor (white-cell growth support) (G-CSF) order is populated for this endocrine monotherapy. Review persistent nausea or other symptoms and prescribe individually if needed.

Medicine-specific support

- Tell the team if joint pain, hot flushes, vaginal dryness or sleep disturbance affects daily life. Bone thinning may cause no early symptoms, so keep the agreed bone-health reviews.
- No routine antiemetic, steroid premedication or G-CSF is required by this endocrine monotherapy. Ask for individual symptom support.
- Ask who books each check, when it is due and who reviews it. Tell the team if an expected appointment has not arrived.

- The kidney and liver advice for exemestane differs from some other aromatase inhibitors: a routine dose reduction is not specified, but organ problems still need clinical review. Tell the pharmacist about all medicines and supplements because some can make exemestane less effective. Do not increase the dose yourself. Confirm menopause and any separate ovarian-suppression plan.
- NCCP no renal/hepatic dose adjustment, haemodialysis expected no adjustment; linked SmPC caution. Strong CYP3A4 inducer reduces exposure: review, no Irish-source automatic 50 mg dose.

Individual plan

Prescriber / date: _____ Pharmacy review: _____

Verify allergies, interactions, relevant organ function, treatment setting and the exact medicine before prescribing.

NCCP 00376 v4 · Monitoring and treatment map

NCCP 00376 · VERSION 4

Exemestane · daily hormone therapy

Take exemestane once daily continuously, preferably after a meal. There is no planned week off. Confirm the prior-treatment and duration plan.

Hormone therapy · selected medicine-specific effects. Full details and sources are in the complete reference.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Joints and muscles


Exemestane

Pain or stiffness can affect activity; report persistent or limiting symptoms.

2 Bone strength

Exemestane

Bone loss may be silent. Your risk assessment and bone-density plan matter.



3 Hot flushes / sleep

Exemestane

Warm flushes and disrupted sleep can be treated; tell the team.

4 Vaginal / sexual symptoms

Exemestane

Dryness or discomfort deserves support. Ask before using hormone-containing treatments.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Report persistent limiting joint pain, new severe bone pain or a possible fracture, jaundice or marked new symptoms. Do not wait for the next scheduled review if you are becoming unwell. Severe chest pain, major breathing difficulty, collapse or stroke symptoms: 112 / 999.

SACTCHECK · CHAPTER 04

Your record and contact card

Exemestane · daily hormone therapy · National Cancer Control Programme (NCCP) 00376

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four recording sheets are reusable review periods, not four prescribed cycles. For injections, use the dates given by your team. For daily tablets, treatment continues unless your prescriber changes the plan.

These patient-language prompts are not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) assessment. Urgent symptoms override the form.

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Joint / muscle pain	☐	☐	☐	
Hot flushes / sleep disruption	☐	☐	☐	
Fatigue	☐	☐	☐	
Vaginal / sexual symptoms	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

Report persistent limiting joint pain, new severe bone pain or a possible fracture, jaundice or marked new symptoms. Do not wait for the next scheduled review if you are becoming unwell. Emergency symptoms: 112 / 999.

Carry your treatment details

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · HORMONE THERAPY

Exemestane · daily hormone therapy

NCCP 00376 v4

Take exemestane once daily continuously, preferably after a meal. There is no planned week off. Confirm the prior-treatment and duration plan.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Mon–Fri 8am–5pm.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Report persistent limiting joint pain, new severe bone pain or a possible fracture, jaundice or marked new symptoms. Do not wait for the next scheduled review if you are becoming unwell.

https://sactcheck.com/patient/00376/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.20 · Sources checked 2026-10-25
Source reconciled; independent clinical, pharmacy and patient/PPI review pending

SACTCHECK · CHAPTER 05

Sources and further information

Exemestane · daily hormone therapy · National Cancer Control Programme (NCCP) 00376

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **NCCP 00376 v4 · official regimen**
   https://healthservice.hse.ie/documents/6568/376_Exemestane_therapy.pdf
2. **HPRA Exemestane SmPC cited by NCCP**
   https://assets.hpra.ie/products/Human/23117/Licence_PA0822-111-001_09012025155108.pdf
3. **Pan-Asian adapted ESMO early breast cancer guidelines**
   https://pmc.ncbi.nlm.nih.gov/articles/PMC11145753/
4. **Pan-Asian adapted ESMO metastatic breast cancer guidelines**
   https://pmc.ncbi.nlm.nih.gov/articles/PMC10186487/
5. **NICE NG 101: early and locally advanced breast cancer**
   https://www.nice.org.uk/guidance/ng101/chapter/Recommendations
6. **ESO-ESMO ABC 5 consensus: tumour-marker interpretation**
   https://pmc.ncbi.nlm.nih.gov/articles/PMC7510449/
7. **IOF/CABS/ECTS/IEG/ESCEO/IMS/SIOG updated joint statement on aromatase-inhibitor-associated bone loss**
   https://www.esceo.org/sites/esceo/files/pdf/1-s2.0-S2212137425000351-main.pdf
8. **Long-term follow-up of the Intergroup Exemestane Study**
   https://pubmed.ncbi.nlm.nih.gov/28467729/
9. **eviQ 3650 v 6: adjuvant whole-breast radiotherapy**
   https://www.eviq.org.au/radiation-oncology/breast/3650-breast-invasive-cancer-adjuvant-whole-breast
10. **Exemestane versus megestrol after tamoxifen failure in advanced breast cancer**
    https://pubmed.ncbi.nlm.nih.gov/10735887/