Trifluridine/tipiracil (Lonsurf) monotherapy

National Cancer Control Programme (NCCP) 00382 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Prescription and supply

Before collecting another course, confirm the dose, tablet strengths, calendar, blood-test date and authorisation. Bring the current packs and medicines list.

STEP 2

Meal routine

Swallow the prescribed tablets with water within one hour after finishing breakfast and the evening meal on dosing days. Doses are based on body size and previous side effects; two tablet strengths may be combined. Some prescriptions have different morning and evening tablet quantities. Use the pharmacy calendar and do not calculate doses yourself.

STEP 3

Missed or vomited doses

Do not make up a missed, held or vomited dose and do not double the next dose. Tell your team and follow the next scheduled dose on the agreed calendar. Repeated vomiting needs advice because of dehydration and missed medicines.

STEP 4

Appointments

Ask who books extra tests, who reviews results and whom to call if the plan is unclear. There is no infusion visit or take-home pump in monotherapy.

STEP 5

Swallowing and handling

Swallow tablets whole with water and wash your hands after handling them. If you cannot swallow tablets whole, ask the pharmacist before changing them. Keep them out of children's reach and ask the pharmacist about safe disposal.

STEP 6

Accidental extra tablets

If you take more Lonsurf than prescribed, contact a doctor or go to hospital straight away. Take the packs and current calendar with you.

Sources: NCCP 00382 v4 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.