Confirm your alectinib treatment setting This guide covers alectinib alone for ALK-positive lung cancer. Confirm whether it is adjuvant treatment after complete surgery, first treatment for advanced cancer, or treatment after crizotinib. These settings have different evidence and treatment lengths.

NATIONAL CANCER CONTROL PROGRAMME (NCCP) 00401 · VERSION 6

Alectinib monotherapy

Take alectinib twice daily with food, every day, with no planned rest week. After surgery, treatment lasts up to 24 months, stopping sooner for recurrence or unacceptable side effects. For advanced cancer, treatment continues until progression or unacceptable side effects under the oncology plan.

BASELINE → TREATMENT → FOLLOW-UP

Treatment and monitoring over time

Intervals are reference guidance, not booked appointments. Ask who arranges tests, reviews results and authorises treatment.

Check	Before starting	First month	Months 2–3	From month 4	Completion / change
Alectinib	Confirm setting and prescription	Twice daily with food	Continue daily if authorised	Continue with review	Adjuvant maximum 24 months; advanced duration depends on benefit/toxicity
Liver tests	Baseline liver profile	Every 2 weeks	Every 2 weeks	Every 4 weeks or as needed	Follow unresolved abnormalities
creatine phosphokinase (CPK) / muscles	Baseline CPK	Every 2 weeks	As clinically indicated	As clinically indicated	Check recovery if held
Blood count / kidneys	Baseline full blood count (FBC) and renal profile	Every 4 weeks	Every 4 weeks	Every 4 weeks	Follow anaemia and renal concerns
Pulse / blood pressure / electrocardiogram (ECG)	Baseline blood pressure (BP) and ECG	Agree early repeat ECG timing	Pulse/BP and ECG as needed	Repeat as clinically indicated	Confirm recovery and medicine review
Cancer and symptoms	Confirm aim and disease assessment	Report new symptoms promptly	Review tolerability and adherence	Imaging per Consultant plan	Agree follow-up or next treatment

THE WHOLE TREATMENT JOURNEY

Alectinib monotherapy: treatment and checks

ALK-positive non-small-cell lung cancer (NSCLC) · adjuvant or advanced disease

Take alectinib twice daily with food, every day, with no planned rest week. After surgery, treatment lasts up to 24 months, stopping sooner for recurrence or unacceptable side effects. For advanced cancer, treatment continues until progression or unacceptable side effects under the oncology plan.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	First month	Months 2–3	From month 4	Completion / change
Alectinib NCCP 00401 · SMPC	Confirm setting and prescription	Twice daily with food	Continue daily if authorised	Continue with review	Adjuvant maximum 24 months; advanced duration depends on benefit/toxicity
Liver tests NCCP 00401 · SMPC	Baseline liver profile	Every 2 weeks	Every 2 weeks	Every 4 weeks or as needed	Follow unresolved abnormalities
CPK / muscles NCCP 00401 · SMPC	Baseline CPK	Every 2 weeks	As clinically indicated	As clinically indicated	Check recovery if held
Blood count / kidneys NCCP 00401 · SMPC	Baseline FBC and renal profile	Every 4 weeks	Every 4 weeks	Every 4 weeks	Follow anaemia and renal concerns
Pulse / blood pressure / ECG NCCP 00401 · SMPC	Baseline BP and ECG	Agree early repeat ECG timing	Pulse/BP and ECG as needed	Repeat as clinically indicated	Confirm recovery and medicine review
Cancer and symptoms NCCP 00401 · SMPC	Confirm aim and disease assessment	Report new symptoms promptly	Review tolerability and adherence	Imaging per Consultant plan	Agree follow-up or next treatment

What each check is for

1 Confirm setting and baseline

Before starting

Sources: NCCP 00401 · SMPC

2 Early liver and muscle checks

First month and first 3 months

Sources: NCCP 00401

3 Ongoing reviews

Throughout treatment

Sources: NCCP 00401 · SMPC

4 Adjuvant completion or advanced change

At planned endpoint or earlier clinical change

Sources: NCCP 00401 · SMPC

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

SACTCheck · Developed by Dr Paul O'Brien · v0.84.24 · Sources checked 2026-10-29

Source reconciled; independent clinical, pharmacy and patient/PPI review pending