Adjuvant dabrafenib and trametinib after removal of BRAF-mutated stage III melanoma

After complete removal of BRAF V600 stage III melanoma; aims to lower recurrence risk.

NCCP 00415b: adjuvant treatment after complete resection of stage III melanoma with a BRAF V600 mutation. This route is distinct from treatment of unresectable or metastatic melanoma. · NCCP 00415 v5 · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

NCCP 00415b: adjuvant treatment after complete resection of stage III melanoma with a BRAF V600 mutation. This route is distinct from treatment of unresectable or metastatic melanoma. This information supports individual consent; it does not issue or change a prescription.

My overall treatment

Both medicines daily; adjuvant course planned for 12 months with regular safety reviews.

28 day view

Treatment timing in this window

Daily oral treatment — Day 1–28

Adjuvant oral treatment

Planned 12 months; no routine drug holiday.

A 28-day reference calendar

Day 1–28

Every day in the reference calendar

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP / medicine-specific precautions	BRAF result; bloods, electrocardiogram (ECG)/ echo, skin exam; eye review if indicated.
During	**Bloods and organ checks** NCCP; local interval clarification where stated	Monthly bloods. ECG/echo at 4 weeks, then 12-weekly and after dose changes. Team records skin checks.
After / response	**Follow-up** Regimen / individual disease plan	Skin surveillance continues 6 months after stopping. Scans and other follow-up as recorded.

Between treatments

- Daily medicines; no routine drug-free week. Fever of 38°C or higher: hold both and contact the team.
- Restart after fever only under the team's plan, after at least 24 hours symptom-free.

Bring the written medicine plan, test dates and actual oncology/out-of-hours contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Fever / chills

Both medicines

At 38°C or above, hold both medicines and contact the team under your fever plan. Infection and dehydration must also be considered.

2 Eyes / vision

Dabrafenib / trametinib

Report blurred vision, eye pain, redness, light sensitivity or a change in sight promptly; some eye problems need urgent specialist treatment.

3 Heart / circulation


Trametinib / combination

Keep ECG and heart ultrasound checks. New breathlessness, swollen ankles, palpitations or chest pain needs prompt assessment.

4 Skin / new lesions

Both medicines

Rash and skin sensitivity may occur; new changing spots need review. Blistering, peeling or mouth sores with a rash needs urgent help.



5 Liver

Both medicines

Liver changes may first appear in blood tests. Report yellow eyes or skin, dark urine, unusual itching or upper-abdominal pain.

6 Kidneys / fluids

Both medicines / fever

Fever, vomiting or diarrhoea can cause dehydration and kidney problems. Call for reduced urine, dizziness or inability to keep fluids down.

7 Muscles / joints

Both medicines

Aches can occur. Severe muscle pain or weakness, especially with dark urine, needs urgent advice and possible blood tests.

8 Stomach / bowels

Both medicines

Tell the team about persistent diarrhoea, vomiting or pain. Severe abdominal pain, blood or black stools needs urgent assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever is a particular concern. At 38°C or above, withhold both medicines and contact the oncology team promptly under the written fever plan; recurrent episodes may require action at the first symptoms. Infection and dehydration also need assessment. Restart only when the team tells you. Urgent help is also needed for vision change, severe rash, chest pain, breathlessness, bleeding or severe muscle symptoms.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

NCCP 00415b: adjuvant treatment after complete resection of stage III melanoma with a BRAF V600 mutation. This route is distinct from treatment of unresectable or metastatic melanoma. · NCCP 00415 v5 · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Dabrafenib + Trametinib

Both medicines daily; adjuvant course planned for 12 months with regular safety reviews.

Phase: _____ Review plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My review record

Use a dated month/review record for the planned 12-month adjuvant course; 12 months does not mean 12 doses or 12 cycles.

Review	Date / initials

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each completed review. Daily tablets continue according to your prescribed plan; a review stamp does not mean the course is complete or authorise further treatment. Record interruptions or a changed plan. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.