National Cancer Control Programme (NCCP) 00430 · clinician draft

Gemcitabine + carboplatin AUC2

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Day 1 · Gemcitabine + carboplatin AUC2

Active medicines: Carboplatin + Gemcitabine

Emetogenic risk: Moderate

Medicine	Dose / route	When
Ondansetron	16 mg orally once	Before chemotherapy on Day 1 Review QT risk, constipation, hepatic function and any alternative local regimen.
Dexamethasone	8 mg orally once daily	Days 1, 2 and 3 Reconcile total steroid exposure; individual adjustments require prescribing review.
Metoclopramide	10 mg orally up to three times daily as needed	Rescue duration to be specified Review renal function, contraindications and interactions before prescribing.

Additional phase-specific support

- Confirm allergy/interaction review, blood tests and the source-specific renal/hepatic assessment before prescribing.
- Carboplatin is AUC2 on BOTH Days 1 and 8; do not use an AUC4/5 Day 1-only template. The AUC≥4 antiemetic category does not apply to this AUC2 schedule.
- Confirm the dose is going ahead before starting the linked support course. Reconcile rescue medicine duration rather than copying another treatment day.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00430 v6 · p5 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.18 · Regimen-level drafts; prescriber and oncology-pharmacy verification required

Day 8 · Gemcitabine + carboplatin AUC2

Active medicines: Carboplatin + Gemcitabine

Emetogenic risk: Moderate

Medicine	Dose / route	When
Ondansetron	16 mg orally once	Before chemotherapy on Day 8 Review QT risk, constipation, hepatic function and any alternative local regimen.
Dexamethasone	8 mg orally once daily	Days 8, 9 and 10 Reconcile total steroid exposure; individual adjustments require prescribing review.
Metoclopramide	10 mg orally up to three times daily as needed	Rescue duration to be specified Review renal function, contraindications and interactions before prescribing.

Additional phase-specific support

- Confirm allergy/interaction review, blood tests and the source-specific renal/hepatic assessment before prescribing.
- Carboplatin is AUC2 on BOTH Days 1 and 8; do not use an AUC4/5 Day 1-only template. The AUC≥4 antiemetic category does not apply to this AUC2 schedule.
- Confirm the dose is going ahead before starting the linked support course. Reconcile rescue medicine duration rather than copying another treatment day.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00430 v6 · p5 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-25 · SACTCheck v0.84.18 · Regimen-level drafts; prescriber and oncology-pharmacy verification required