FOLFIRI + bevacizumab

FOLFIRI + bevacizumab is used for cancer of the colon or rectum that has spread. Treatment aims to control the cancer; the likely benefit and review plan depend on your individual situation. Surgery may be reconsidered in selected cases after multidisciplinary review.

Metastatic colorectal cancer · NCCP 00449 v6a · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Day 1: bevacizumab, irinotecan, folinic acid and fluorouracil, then a 46-hour fluorouracil pump. A cycle lasts 14 days; your team confirms the next treatment.



14-day treatment cycle

Day 1: bevacizumab, irinotecan, folinic acid and fluorouracil, then a 46-hour fluorouracil pump. A cycle lasts 14 days; your team confirms the next treatment.

Your treatment schedule

Day 1	Days 2–3	Days 4–14
Day-ward medicines; pump connected.	Complete 46-hour pump; disconnect as arranged.	Recovery, symptoms and tests; next cycle only after team review.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**dihydropyrimidine dehydrogenase (a medicine-processing enzyme) (DPD) / baseline bloods** Required by regimen	DPD before fluoropyrimidine; blood count, kidney and liver tests.
During	**Bloods** Required by regimen	Blood count, kidney and liver tests before each cycle.
Before / during	**blood pressure (BP) / urine protein** Required by regimen	Baseline and each-cycle urine protein; BP before each cycle and after treatment.
If indicated	**Heart / international normalised ratio (blood-clotting test) (INR)** Conditional tests	electrocardiogram (ECG) if cardiac function is compromised. Warfarin: INR weekly until stable, then before each cycle.
After / review	**Recovery / scans / markers** Team-agreed plan	Follow the individual cancer plan and any unresolved effects; scan and marker intervals are team-agreed.

Between treatments

- No guaranteed safe days: report fever, sudden illness, diarrhoea or poor intake promptly.
- Symptoms can continue after pump removal and after treatment ends.

Confirm supportive medicines, tests, working urgent contacts and pump disconnection.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hair

Irinotecan

Hair thinning or loss may occur. Ask about support.

2 Mouth

Fluorouracil + folinic acid

Report ulcers or pain affecting eating and drinking.

3 Chest

Fluorouracil / bevacizumab

New chest pain or breathlessness needs urgent assessment.

4 Bowel

Irinotecan / fluorouracil

Report diarrhoea, increased stoma output or severe abdominal pain.

5 Blood pressure / kidneys

Bevacizumab

Keep BP and urine checks. Report severe headache or reduced urine.

6 Blood / infection / bleeding

Chemotherapy / bevacizumab

Fever or sudden illness: call now. Major bleeding or a painful swollen limb needs urgent review.

7 Skin / hands / feet


Fluorouracil

Report painful skin changes affecting activity.

8 Pump / line / wounds

Treatment delivery / bevacizumab

Pain, leakage, swelling or poor wound healing needs advice.



Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact promptly for new chest symptoms, severe or persistent diarrhoea, reduced urine, inability to drink, severe abdominal pain or a line/pump problem. New neurological symptoms, uncontrolled bleeding or a swollen painful limb also need urgent assessment.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Metastatic colorectal cancer · NCCP 00449 v6a · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

FOLFIRI + bevacizumab

Every 14 days · 46-hour fluorouracil pump · cycle plan confirmed by team.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.