National Cancer Control Programme (NCCP) 00507 · clinician draft

Weekly paclitaxel + pertuzumab + trastuzumab

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Days 8 and 15 · paclitaxel only

Active medicines: Paclitaxel

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- Day 8 and later: no routine dexamethasone unless consultant-directed for reaction/risk; chlorphenamine continues, and famotidine omission requires consultant/local-policy agreement.
- Reconcile antiemetic steroids with taxane premedication; do not duplicate.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.
- No HER2 antibodies on these weekly-only visits.
- Clinical template for individual review, not an issued prescription. Confirm allergy, pregnancy precautions, interactions and local policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00507 v6 · p5 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-21 · SACTCheck v0.84.16 · Regimen-level drafts; prescriber and oncology-pharmacy verification required