INFORMATION TO KEEP WITH YOU

My complete treatment handbook

Ribociclib + endocrine therapy · metastatic

Understand your treatment. Know what to expect. Keep your plan to hand.

Displayed calendar
28 days
See the active treatment phase

Treatment stages
1
Explained in the treatment reference

Regimen source
NCCP 00525
Version 6

Oral treatment with a planned rest week

Treatment stages

01 **A 28-day reference block**
Treatment → checks → review

Your team confirms the active stage and dates. The full treatment guide explains each step.

Inside your handbook

		PAGE
01	Understanding your treatment	2
02	Your schedule, tests and appointments	4
03	Symptoms and supportive care	7
04	Your record and contact card	12
05	Sources and further information	14

My name _____

Hospital / oncology team _____

Use the urgent instructions and contact numbers in your treatment guide. Your team must replace example numbers before patient use. Your individual treatment plan takes precedence.

SACTCHECK · CHAPTER 01

Understanding your treatment

Ribociclib + endocrine therapy · metastatic · National Cancer Control Programme (NCCP) 00525

How this medicine supports your treatment

Ribociclib with an aromatase inhibitor or fulvestrant aims to control advanced HR-positive/HER2-negative breast cancer. The exact endocrine partner depends on previous treatment and menopausal status.

CDK4/6 INHIBITOR

Ribociclib

Slows the cell-division checkpoint

CDK4/6 checkpoint → Checkpoint inhibited → Cell cycling slows

Ribociclib blocks CDK4 and CDK6, proteins that help cells move through the cycle of division. Endocrine therapy reduces the hormone signal that can drive HR-positive breast cancer.

Keep blood and electrocardiogram (ECG) appointments. Fever, jaundice, fainting, palpitations, severe rash or new breathlessness need prompt advice.

ENDOCRINE PARTNER

aromatase inhibitor (AI) or fulvestrant

Reduces oestrogen-driven growth

Oestrogen pathway → AI or receptor blockade → Growth signal reduced

An aromatase inhibitor (AI) lowers oestrogen production; fulvestrant blocks and reduces oestrogen receptors. These are alternative endocrine partners, selected by the team. Ovarian suppression is needed in pre/perimenopausal patients.

The endocrine medicine continues during the ribociclib week off. Report troublesome joint, menopausal or injection-site symptoms.

Your treatment aim

Who this treatment is for

HR-positive/HER2-negative locally advanced or metastatic disease with an AI or fulvestrant in the exact NCCP indication. NCCP page 1 and page 2 do not fully align on menopause and previous advanced-disease treatment; preserve consultant reconciliation.

NCCP 00525 v6 · official regimen

Schedule and duration

The usual metastatic starting dose is ribociclib 600 mg once daily on Days 1–21 followed by Days 22–28 off. Continue the prescribed AI daily or fulvestrant on its own injection schedule; add ovarian suppression when indicated. Continue while beneficial and tolerated. There is no fixed three-year stop in this metastatic regimen; the team reviews response, side effects and the next treatment decision.

NCCP 00525 v6 · official regimen

What the studies tell us

Studies of ribociclib with hormone therapy in defined advanced-breast-cancer groups showed a survival benefit. The studies used different hormone partners and included different menopausal and prior-treatment groups. Ask which study matches your plan; the results do not set an individual time limit.

How to interpret this

Study results describe groups with defined treatment and eligibility. Ask about your own likely benefits, important harms, alternatives and remaining uncertainty.

MONALEESA-2 final OS, Hortobagyi et al 2022 · MONALEESA-2 updated PFS, Hortobagyi et al 2018 · MONALEESA-3 updated overall-population OS, Slamon et al 2021 · MONALEESA-7 exploratory extended OS, Lu et al 2022

MONALEESA-2, MONALEESA-3 and MONALEESA-7: separate populations and partners

Studies describe groups. Ask how closely the treatment and study population match your situation.

Your team can discuss the strength and limitations of the evidence alongside treatment risks and alternatives.

When to get help

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Report new breathlessness, chest symptoms, severe diarrhoea, inability to drink, jaundice, a painful swollen leg or a severe rash promptly. You do not need to identify which medicine caused the symptom.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

SACTCHECK · CHAPTER 02

Your schedule, tests and appointments

Ribociclib + endocrine therapy · metastatic · NCCP 00525

BASELINE → TREATMENT → FOLLOW-UP

Treatment and monitoring over time

Columns show sequence and general intervals, not equal periods or booked appointments. The team confirms who arranges each test, who reviews results and who authorises treatment or restart.


The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00525 v6 · official regimen · EMA Kisqali current product information; checked 2026-10-25

THE WHOLE TREATMENT JOURNEY

Ribociclib + endocrine therapy · metastatic: treatment and checks

Ribociclib + endocrine therapy · metastatic

28-day view



This shows treatment timing, not guaranteed symptom-free days. Your team confirms the active phase and dates.

HR-positive / HER2-negative metastatic breast cancer

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before first dose	Cycles 1–2	Cycles 3–6	Later treatment	Completion / change
Ribociclib and endocrine partner NCCP 00525 · SMPC	Confirm metastatic 600 mg / AI or fulvestrant plan	Days 1–21 on; Days 22–28 off; fulvestrant loading visits if chosen	Repeat if cleared; endocrine partner continues	While beneficial and tolerated	Review endocrine continuation separately
Blood counts / renal / liver NCCP 00525 · SMPC	Baseline profiles	Every two weeks in both cycles	Before each cycle	As clinically indicated; more often if abnormal	Follow abnormalities before restart or after stopping

Care / check	Before first dose	Cycles 1–2	Cycles 3–6	Later treatment	Completion / change
ECG / heart rhythm NCCP 00525 · SMPC	Baseline ECG; check QTcF eligibility	Repeat around cycle 1 Day 14	Further ECG when clinically indicated	Repeat for symptoms, QT change or interacting medicines	No automatic restart after a QT-related hold
Electrolytes NCCP 00525 · SMPC	Potassium, calcium, magnesium and phosphorus	Before each cycle; correct abnormalities	Before each cycle through cycle 6	As clinically indicated	Review unresolved abnormalities
Bone / endocrine health NCCP 00525 · SMPC	Assess AI/ ovarian function suppression (OFS) bone and menopausal risks	Support symptoms and adherence	AI/OFS and any injections follow their own plan	Repeat bone assessment according to risk	Longer endocrine and bone plan continues
Cancer response NCCP 00525 · SMPC	Baseline disease assessment	Assess new symptoms promptly	Imaging according to consultant plan	Continue response and symptom review	Progression prompts new treatment decision

What each check is for

1 Before the first dose

Before first dose

Check eligibility, organ function and the exact treatment/support plan.

Ask / act: Confirm who arranges each check and who reviews it.

2 Early toxicity checks

Cycles 1–2

Important toxicity can begin before the first routine review.

Ask / act: Keep early appointments and use the agreed symptom-contact plan.

3 Benefit, tolerability and adherence

Later treatment

Longer-term treatment still needs clinical review.

Ask / act: Ask about symptom support, results and the next decision.

4 Completion or treatment change

Completion / change

The targeted medicine and endocrine partner may have different stop dates.

Ask / act: Confirm each medicine and the ongoing follow-up plan.

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Checks before and during treatment

Baseline tests and ongoing tests have different purposes.

Before the first dose

full blood count (FBC), renal and liver profile, potassium, calcium, magnesium and phosphorus. ECG before initiation; QTcF must be below 450 ms. Correct electrolyte abnormalities and review QT/CYP3A interactions. Confirm endocrine partner, menopause/ovarian suppression, prior treatment and the NCCP indication/eligibility discrepancy.

During treatment

FBC, renal and liver profile every two weeks for cycles 1–2, then before cycles 3–6, then as clinically indicated. Electrolytes before each of the first six cycles, then as clinically indicated; correct abnormalities. Repeat ECG around Day 14 of cycle 1, then as clinically indicated. Increase liver monitoring for grade 2 or worse abnormalities. Review fever, jaundice, palpitations/fainting, pulmonary and severe skin symptoms; response imaging follows the advanced-disease plan.

Practical treatment details

Your endocrine partner has a separate timetable

If fulvestrant is the chosen partner, the usual schedule is 500 mg by intramuscular injection on Days 1, 15 and 29, then once monthly; confirm the prescribed dose and appointment dates.

Supply and appointments

Request the next supply before tablets run out.

This oral treatment does not itself require a central line or infusion pump.

Support between appointments

Keep the oncology contact details and prescribed symptom medicines accessible.

SACTCHECK · CHAPTER 03

Symptoms and supportive care

Ribociclib + endocrine therapy · metastatic · NCCP 00525

Day 1 → Day 28: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Treatment or check	Timing in this view
Ribociclib tablets	 **Days 1–21** Once daily
Ribociclib planned break	 **Days 22–28** No ribociclib
Endocrine partner	 **Days 1–28** AI daily OR injection schedule
Blood counts / infection	 **Days 1–28** Tests and symptoms guide review
Energy and bowel symptoms	 **Days 1–28** Timing varies; report changes early
Heart rhythm / ECG	 **Days 1–28** Day-14 ECG in cycle 1; symptoms any day
Optional local pre-cycle review	 **Days 25–26** Local D25–26 window

Treatment days: Ribociclib is taken once daily on Days 1–21, followed by seven days off, in a 28-day cycle. The endocrine partner continues according to its own prescription. Treatment continues while it is helping and tolerable.

The bigger picture: Use the whole-treatment journey for surgery, radiation, scans and longer-term checks.

Read the phase labels: The prescription and team's instructions determine your actual doses and any interruptions.

Local pre-cycle review: Day 25–26 can be a local blood / review window. It does not replace the required early checks or authorise an automatic restart.

Why these bands stay broad: Treatment bands show the source schedule. Symptom bands show possible patterns, not measured probabilities; serious effects can occur at any time. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00525 v6 · official regimen ↗

There are no guaranteed "safe days". Fever, shivers or suddenly feeling unwell require immediate contact. Serious symptoms override the timeline or diary.

Living with treatment

Supportive medicines and the endocrine partner have their own plan.

Nausea support

NCCP classifies ribociclib as oral moderate–high emetic risk. The team chooses preventive or rescue antiemetics using local policy, symptoms, QT risk and drug interactions; do not reuse another regimen's prescription.

Electrolytes and heart rhythm

Correct abnormal potassium, calcium, magnesium and phosphate as directed. Attend the baseline and cycle 1 Day-14 ECG; fainting or palpitations need urgent advice.

Counts and liver tests

Keep the early two-weekly blood tests and subsequent monitoring. Report fever, shivers, jaundice or dark urine promptly.

Endocrine treatment

The AI or fulvestrant, and ovarian suppression when prescribed, continue on their own timetable. Discuss joint, menopausal, bone and injection-site symptoms.

Daily administration

Avoid grapefruit and check interactions. Do not replace a missed or vomited dose. A clinical hold requires explicit instructions about which medicine stops and who authorises restart.

When the plan changes

Continue, hold or stop

The team confirms the plan for each medicine and the next review. Do not restart a clinically held treatment from this educational timeline.

Practical support

Nausea support

Electrolytes and heart rhythm

Counts and liver tests

Endocrine treatment

Daily administration

Taking tablets / attending treatment

Questions to ask

Which checks happen before and during treatment?

FBC, renal and liver profile, potassium, calcium, magnesium and phosphorus. ECG before initiation; QTcF must be below 450 ms. Correct electrolyte abnormalities and review QT/CYP3A interactions. Confirm endocrine partner, menopause/ovarian suppression, prior treatment and the NCCP indication/eligibility discrepancy. FBC, renal and liver profile every two weeks for cycles 1–2, then before cycles 3–6, then as clinically indicated. Electrolytes before each of the first six cycles, then as clinically indicated; correct abnormalities. Repeat ECG around Day 14 of cycle 1, then as clinically indicated. Increase liver monitoring for grade 2 or worse abnormalities. Review fever, jaundice, palpitations/fainting, pulmonary and severe skin symptoms; response imaging follows the advanced-disease plan.

Do I need to enter dates?

No. This map shows general timing without personal information. Your hospital arranges actual appointments; ask who books each test and who reviews its result.

What if a dose is missed or treatment is held?

Take ribociclib once daily with or without food; swallow whole. Avoid grapefruit. If a dose is missed or vomited, do not take an extra dose; take the next scheduled dose. Confirm any clinical hold and restart with the team. The endocrine partner follows a separate prescription. A hold may affect one medicine rather than the whole combination; confirm each component.

What happens after treatment changes or ends?

There is no fixed three-year stop in this metastatic regimen; the team reviews response, side effects and the next treatment decision. Keep the separately agreed disease follow-up, endocrine and bone-health plan.

Explain the plan back: Which medicine is taken when, which test is due next, who reviews it, and which symptoms mean you should call today?

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Ribociclib + endocrine therapy · metastatic · supportive-care plan

Active medicines: Ribociclib + aromatase inhibitor (AI) or fulvestrant

Emetogenic risk: Oral moderate–high

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- NCCP classifies ribociclib as oral moderate–high emetic risk. The team chooses preventive or rescue antiemetics using local policy, symptoms, QT risk and drug interactions; do not reuse another regimen's prescription.
- Correct abnormal potassium, calcium, magnesium and phosphate as directed. Attend the baseline and cycle 1 Day-14 electrocardiogram (ECG); fainting or palpitations need urgent advice.
- Keep the early two-weekly blood tests and subsequent monitoring. Report fever, shivers, jaundice or dark urine promptly.
- The AI or fulvestrant, and ovarian suppression when prescribed, continue on their own timetable. Discuss joint, menopausal, bone and injection-site symptoms.
- Avoid grapefruit and check interactions. Do not replace a missed or vomited dose. A clinical hold requires explicit instructions about which medicine stops and who authorises restart.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Heart rhythm / QT

Ribociclib

Baseline ECG and an ECG around Cycle 1 Day 14. Fainting or palpitations need urgent advice.

3 Low neutrophils / infection


Ribociclib

Blood counts guide whether the next cycle can start. Fever, shivers or sudden illness: contact oncology immediately.

5 Severe rash or blistering

Ribociclib

A widespread painful rash, blisters or peeling skin needs urgent assessment.



2 Liver injury

Ribociclib

Keep liver blood tests. Yellow skin / eyes, dark urine or marked upper-abdominal discomfort: call promptly.

4 Lung inflammation

Ribociclib · uncommon but important

New cough or breathlessness needs prompt assessment, even between visits.

6 Joints and bone health

Aromatase inhibitor / ovarian suppression

Discuss joint symptoms and keep the agreed bone-risk and DXA plan.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Report new breathlessness, chest symptoms, severe diarrhoea, inability to drink, jaundice, a painful swollen leg or a severe rash promptly. You do not need to identify which medicine caused the symptom.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

SACTCHECK · CHAPTER 04

Your record and contact card

Ribociclib + endocrine therapy · metastatic · National Cancer Control Programme (NCCP) 00525

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four sheets are reusable review periods, not a prescribed number of cycles. Follow the treatment and rest days agreed with your team; do not calculate a restart from this sheet.

These patient-language prompts are not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) assessment. Urgent symptoms override the form.

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Nausea	☐	☐	☐	
Fatigue	☐	☐	☐	
Skin / nail changes	☐	☐	☐	
Muscle / joint pain	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

Report new breathlessness, chest symptoms, severe diarrhoea, inability to drink, jaundice, a painful swollen leg or a severe rash promptly. You do not need to identify which medicine caused the symptom. Emergency symptoms: 112 / 999.

Keep your treatment details to hand

HR-positive / HER2-negative metastatic breast cancer · NCCP 00525 v6 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Ribociclib + endocrine therapy

Ribociclib Days 1–21 of 28; endocrine partner has a separate continuing schedule.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.32 · NCCP 00525 v6 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Sources and further information

Ribociclib + endocrine therapy · metastatic · National Cancer Control Programme (NCCP) 00525

Use this information alongside the plan agreed with your treatment team. Your team confirms your medicines, doses and appointments.

1. **NCCP 00525 v6 · official regimen ↗**
   https://healthservice.hse.ie/documents/6629/525_v6_Ribociclib_Metastatic_hghzyyD.pdf
2. **EMA Kisqali current product information; checked 2026-10-25**
   https://www.ema.europa.eu/en/documents/product-information/kisqali-epar-product-information_en.pdf
3. **NICE NG101 · bone health and follow-up**
   https://www.nice.org.uk/guidance/ng101/chapter/Recommendations
4. **HSE · chemotherapy side effects and urgent advice**
   https://www2.hse.ie/conditions/chemotherapy/side-effects/
5. **ASCO · biomarkers in metastatic breast cancer**
   https://ascopubs.org/doi/10.1200/JCO.2015.61.1459
6. **ESMO metastatic breast cancer guideline · response assessment**
   https://www.annalsofoncology.org/article/S0923-7534(26)00902-6/fulltext
7. **NOGG · section 7, aromatase inhibitors and fracture risk**
   https://www.nogg.org.uk/full-guideline/section-7-strategies-management-osteoporosis-and-fracture-risk
8. **eviQ · peripherally inserted central catheter (PICC) patient information**
   https://www.eviq.org.au/p/543
9. **eviQ · implanted port patient information**
   https://www.eviq.org.au/p/507
10. **Novartis · Kisqali product information, updated September 2026**
    https://www.medicines.org.uk/emc/product/8110/smpc
11. **MONALEESA-2: ribociclib with letrozole · primary report**
    https://pubmed.ncbi.nlm.nih.gov/35263519/
12. **MONALEESA-2 updated PFS, Hortobagyi et al 2018**
    https://pubmed.ncbi.nlm.nih.gov/29718092/
13. **MONALEESA-3 updated overall-population OS, Slamon et al 2021**
    https://pubmed.ncbi.nlm.nih.gov/34102253/
14. **MONALEESA-7 exploratory extended OS, Lu et al 2022**
    https://pubmed.ncbi.nlm.nih.gov/34965945/
15. **NCCP antiemetics v8, Table 1 pp4–5**
    https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf