
# Pertuzumab + trastuzumab + vinorelbine

intravenous (IV) pertuzumab and IV trastuzumab on Day 1; vinorelbine on Days 1 and 8 of each 21-day cycle. Vinorelbine is prescribed by IV OR by mouth, not both. There is no planned Day 15 vinorelbine. Continue while beneficial and tolerated; treatment may change if one component causes toxicity.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.



### 1 Heart / breathing

**HER2 antibodies**

Keep echoes; report breathlessness, swelling or chest symptoms.

### 2 Constipation / bowel pain

**Vinorelbine**

Report constipation early; severe pain, vomiting or inability to pass stool or wind needs urgent advice.

### 3 Blood / infection

**Vinorelbine**

Fever, shivers or sudden illness: contact oncology immediately.

### 4 Nerves

**Vinorelbine**

Report tingling, weakness or difficulty with usual tasks.

### 5 Infusion reaction

**HER2 antibodies**

Tell staff immediately about rash, chest tightness or breathing difficulty.

### 6 IV site / oral nausea

**Vinorelbine**

IV burning or swelling needs immediate attention. Oral nausea needs a route-specific plan.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

The map shows selected effects of the medicines named here. Report persistent effects from earlier treatment too.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

# Your treatment, at home.

## Important medicine-specific effects

### Pertuzumab

- Heart pumping-function reduction
- Treatment reaction
- Diarrhoea and rash
- Fatigue

### Trastuzumab

- Heart pumping-function reduction
- Treatment reaction
- Diarrhoea and rash
- Fatigue

### Vinorelbine

- Low blood counts and infection
- Constipation or bowel obstruction
- Numbness or weakness
- Nausea; oral doses need a specific plan
- Irritation if an IV dose leaks

**Surgery, radiation and monitoring:** open the whole-treatment map or scan the QR.

## Make the plan practical

### Route-specific support

Oral vinorelbine needs a separate nausea plan; do not copy an IV-only minimal-risk template. Confirm anti-sickness treatment, constipation prevention and blood checks before each prescribed dose.

### Tests before treatment

full blood count (FBC), kidney and liver profile at baseline and before every 21-day cycle. The team reviews the necessary results before a planned dose. Confirm the blood-count check needed before each additional vinorelbine or paclitaxel dose; a Day 1 result does not automatically authorise a later dose. For oral vinorelbine, the product information requires blood counts on the day of each administration, including Day 8; the regimen source lists full blood/renal/liver checks before each cycle.

### Keep heart checks on the plan

An echo every 12 weeks is separate from blood tests or cancer imaging. Ask who books and reviews it.

### Plan for symptoms

Keep your agreed contact numbers and instructions. New symptoms can need assessment before a routine appointment.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

**Complete reference, visual timeline and passport**

https://sactcheck.com/patient/00526/

Sources and review status are available online. Follow your own prescription and hospital instructions.