Pertuzumab + trastuzumab + vinorelbine

National Cancer Control Programme (NCCP) 00526 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

intravenous (IV) pertuzumab and IV trastuzumab on Day 1; vinorelbine on Days 1 and 8 of each 21-day cycle. Vinorelbine is prescribed by IV OR by mouth, not both. There is no planned Day 15 vinorelbine. Continue while beneficial and tolerated; treatment may change if one component causes toxicity.

STEP 1

Before the first visit

Confirm the active regimen, blood results, cardiac checks, medicines review, route and consent discussion.

STEP 2

Plan venous access

A cannula, peripherally inserted central catheter (PICC) or implanted port may be used. If a port is chosen, a trained team inserts it in a separate procedure with local anaesthetic and sometimes sedation; the team explains fasting, blood-thinner and wound instructions.

STEP 3

Infusion day

The HER2 antibodies still require IV access even when vinorelbine is taken orally. Vinorelbine is given or taken on Days 1 and 8; the loading-antibody visit takes longer. A Day 8 plan may include blood review and tablets rather than an infusion, as arranged by your team.

STEP 4

Home and next review

This regimen does not routinely need a take-home infusion pump. Staff disconnect the infusion; a port is de-accessed when appropriate. Keep line-care and next-visit instructions.

Port insertion and removal

The line team explains placement, dressing care and when normal activities can resume. A port is accessed with a special needle during treatment. Removal of the implanted device is a separate planned procedure when no longer needed; do not remove or adjust it yourself.

PICC care and urgent line symptoms

A PICC usually needs regular dressing and flushing appointments even between treatments. Fever, redness, swelling, pain, leakage or damage requires prompt advice. Your hospital supplies the exact care instructions.

Visit time and transport

First visits can take longer because of loading doses, premedicines and observation. Ask the unit about expected time, transport and any escort needed after sedation or line insertion.

Phase-specific visit pattern

Infusion and observation times differ by drug and first versus later doses; the unit confirms the full visit duration, including blood review and preparation.

If taking capsules

Oral vinorelbine is taken only on the prescribed Days 1 and 8, with some food. Swallow capsules whole with water; do not chew, suck, open or dissolve them. Do not use a damaged capsule. Do not take an extra dose after vomiting; contact the team. Dose increases require a new prescription and blood-count review.

Sources: NCCP 00526 v5 · official regimen · eviQ · implanted port patient information · eviQ · PICC patient information. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.