Which treatment this guide covers National Cancer Control Programme (NCCP) 00565a covers first-line treatment of adults with locally advanced or metastatic non-small-cell lung cancer (NSCLC) with an activating EGFR mutation. This is daily dacomitinib alone; it is not an adjuvant regimen or a chemotherapy combination.

NCCP 00565 · VERSION 3

Dacomitinib monotherapy

The usual starting dose is 45 mg by mouth once daily, continuously without a planned rest week. Use the dose on your prescription; some patients start lower because of liver function. Treatment continues while it helps and side effects remain acceptable.

Your treatment and review passport

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four sheets are reusable review periods, not a prescribed number of cycles. Follow the treatment and rest days agreed with your team; do not calculate a restart from this sheet.

These patient-language prompts are not a validated questionnaire or a formal Common Terminology Criteria for Adverse Events (CTCAE) assessment. Urgent symptoms override the form.

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or peeling skin reactions; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit's temperature and 24-hour contact instructions.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

Your complete treatment reference

https://sactcheck.com/patient/00565/

Public information only. No personal information is encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Dacomitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Skin / nail changes	□	□	□	
Mouth soreness / ulcers	□	□	□	
New cough or breathlessness	□	□	□	
Eye pain or blurred vision	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or peeling skin reactions; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit's temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.

Dacomitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Skin / nail changes	□	□	□	
Mouth soreness / ulcers	□	□	□	
New cough or breathlessness	□	□	□	
Eye pain or blurred vision	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or peeling skin reactions; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.

Dacomitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Skin / nail changes	□	□	□	
Mouth soreness / ulcers	□	□	□	
New cough or breathlessness	□	□	□	
Eye pain or blurred vision	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or peeling skin reactions; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.

Dacomitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	□	□	□	
Skin / nail changes	□	□	□	
Mouth soreness / ulcers	□	□	□	
New cough or breathlessness	□	□	□	
Eye pain or blurred vision	□	□	□	
Fatigue	□	□	□	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or peeling skin reactions; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit's temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.