National Cancer Control Programme (NCCP) 00605 · clinician draft

Talazoparib · breast monotherapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Continuous oral treatment

Active medicines: Talazoparib

Emetogenic risk: Oral minimal-to-low

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No routine antiemetic premedication is required. Select a PRN antiemetic under the oral Table 5/local plan; review renal function, QT risk and interactions.
- Monitor blood counts and organ function every four weeks; transfusion or growth-factor support is individualised.
- Review capsule strength, adherence and interacting medicines before issuing a new prescription.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

National Cancer Control Programme (NCCP) 00605 v2 · p4 · NCCP oral antiemetic guidance v6, Tables 5 and 8

NCCP oral antiemetic classification v6 · Tables 5 and 8 · Sources checked 2026-10-25 · SACTCheck v0.84.19 · Regimen-level drafts; prescriber and oncology-pharmacy verification required