Paclitaxel Days 1, 8 and 15 · relapsed or refractory small-cell lung cancer (SCLC)

This treatment aims to control small-cell lung cancer that has returned or has not responded to previous treatment, and may help symptoms.

Your treatment plan

One infusion each week for three weeks, then one week without treatment. This 28-day cycle repeats while helping and tolerated.

1. **Before starting**
   Agree the treatment aim, alternatives, blood tests and support medicines.

2. **Three treatment visits**
   Paclitaxel on Days 1, 8 and 15, with blood tests and symptom review before each dose.

3. **One week off**
   No Day 22 infusion. Keep watching for symptoms and confirm the next cycle.

Before and between visits

- Bring your medicine list and symptom record.
- Report tingling, numbness or weakness before each dose.
- Confirm premedication, blood tests and urgent contact numbers.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately for temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions.

Cannot reach the team? Seek urgent medical assessment. Do not wait for the next appointment.

Your contact plan

Ask your team to complete these before you leave.

Hospital / team _____

Daytime number and hours _____

Evening / weekend number _____

If unavailable, where to go _____

Keep this with your consent information. Ask your team about anything unclear.

What to watch for

Use the picture and symptom descriptions together. You do not need to know which medicine caused a symptom before calling.

1. Hands / feet

Paclitaxel

Report numbness, weakness or changes in walking and hand use before the next dose.

2. Blood / infection

Paclitaxel

Fever, shivers or sudden illness need immediate contact.

3. Hair / scalp

Paclitaxel

Hair loss can occur; ask about support before treatment.

4. Breathing


Paclitaxel

New or worsening breathlessness or cough needs urgent assessment.

5. Mouth

Paclitaxel

Report painful ulcers, difficulty drinking or poor intake.



Selected effects, not every possible effect or a prediction. Match each number to its symptom box. Blood, skin and nerve markers are schematic.

Call the oncology contact number immediately for fever, shivers or sudden illness, severe or new breathing difficulty, chest pain, fainting, severe abdominal pain or persistent diarrhoea. Tell staff immediately about infusion symptoms. Report new numbness, weakness, painful skin changes, swelling or visual changes promptly; do not wait for the next cycle.

My systemic anticancer therapy (SACT) card

Paclitaxel Days 1, 8 and 15 · relapsed or refractory small-cell lung cancer (SCLC)

Paclitaxel Days 1, 8 and 15 every 28 days; no Day 22 dose.

Receiving paclitaxel. Fever or sudden illness needs urgent oncology advice.

Name: _____
Hospital: _____
Oncology phone: _____
Out of hours: _____

Show this card when seeking care. It does not replace a medicine-specific alert card.

More information, when you want it

Scan for medicines, tests, daily life and your treatment record.

https://sactcheck.com/patient/00621/lung/