Adult treatment scope · National Cancer Control Programme (NCCP) 00702a / 00702b This adult suite covers entrectinib alone for ROS1-positive advanced non-small-cell lung cancer (NSCLC) before any ROS1 inhibitor (00702a), or a separate NTRK gene-fusion solid-tumour indication with no previous NTRK inhibitor and no satisfactory treatment options (00702b). Molecular tests and Irish access criteria must be confirmed. The adult 600 mg schedule does not apply to children.

NCCP 00702 · VERSION 2

Entrectinib monotherapy · adult

The usual adult starting dose is 600 mg by mouth once daily, every day without a planned rest week. Use your prescribed dose. Treatment continues while helping and side effects remain acceptable; there is no fixed number of cycles.

BASELINE → TREATMENT → FOLLOW-UP

Treatment and checks over time

Relative milestones, not booked appointments or equal time periods. electrocardiogram (ECG)/electrolytes are checked after one month; daily dosing has no planned rest week.

Check	Before treatment	First month	Ongoing daily treatment	Benefit / toxicity review	Hold or change
Treatment	Confirm ROS1 or NTRK setting and prescription	Daily capsules without a rest week	Continue prescribed dose if authorised	Review benefit and tolerability	Team authorises hold, restart, switch or stop
Blood tests	Blood count, kidney/liver tests and uric acid	Repeat at agreed reviews	Continue regular testing	Extra tests for symptoms/abnormal results	Follow recovery and unresolved effects
Heart checks	ECG/electrolytes; left ventricular ejection fraction (heart pumping measurement) (LVEF) if symptoms/ risk	ECG/electrolytes after one month	Further checks periodically as indicated	Assess new breathlessness/swelling/ fainting	Confirm recovery before restarting
Symptoms and supply	Cognition, vision, neuropathy and medicine list	Review symptoms, adherence and supply	Report thinking/memory, visual or bone changes	Review interactions and practical support	Continue contact and reproductive precautions
Cancer assessment	Validated fusion, stage and prior therapy	Clinical review	Scans at Consultant-directed intervals	Discuss response/progression and preferences	Agree next treatment or follow-up

THE WHOLE TREATMENT JOURNEY

Entrectinib monotherapy · adult: treatment and checks

ROS1-positive advanced NSCLC or separate eligible NTRK fusion-positive tumour

The usual adult starting dose is 600 mg by mouth once daily, every day without a planned rest week. Use your prescribed dose. Treatment continues while helping and side effects remain acceptable; there is no fixed number of cycles.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before treatment	First month	Ongoing daily treatment	Benefit / toxicity review	Hold or change
Treatment NCCP 00702 · SMPC	Confirm ROS1 or NTRK setting and prescription	Daily capsules without a rest week	Continue prescribed dose if authorised	Review benefit and tolerability	Team authorises hold, restart, switch or stop
Blood tests NCCP 00702 · SMPC	Blood count, kidney/liver tests and uric acid	Repeat at agreed reviews	Continue regular testing	Extra tests for symptoms/abnormal results	Follow recovery and unresolved effects
Heart checks NCCP 00702 · SMPC	ECG/electrolytes; LVEF if symptoms/ risk	ECG/electrolytes after one month	Further checks periodically as indicated	Assess new breathlessness/swelling/ fainting	Confirm recovery before restarting
Symptoms and supply NCCP 00702 · SMPC	Cognition, vision, neuropathy and medicine list	Review symptoms, adherence and supply	Report thinking/memory, visual or bone changes	Review interactions and practical support	Continue contact and reproductive precautions
Cancer assessment NCCP 00702 · SMPC	Validated fusion, stage and prior therapy	Clinical review	Scans at Consultant-directed intervals	Discuss response/progression and preferences	Agree next treatment or follow-up

What each check is for

1 Confirm the setting

Before starting

The fusion and previous inhibitor change eligibility.

Ask / **act:** Confirm validated testing, MAP approval, prescription and baseline checks.

Sources: NCCP 00702 · SMPC · MAP

2 One-month heart review

After one month

QT and electrolytes can change without symptoms.

Ask / **act:** Agree who books the ECG/blood tests and reviews the results.

Sources: NCCP 00702 · SMPC

3 Ongoing review

During daily therapy

Benefit, symptoms and organ function can change.

Ask / **act:** Keep the medicine list current and report symptoms between visits.

Sources: NCCP 00702 · SMPC

4 Hold or change

When toxicity/progression requires

Recovery and restart differ by adverse effect.

Ask / **act:** Follow the written clinician plan; continue follow-up and contraception advice.

Sources: NCCP 00702 · SMPC

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

SACTCheck · Developed by Dr Paul O'Brien · v0.84.25 · Sources checked 2026-10-29

Source reconciled; independent clinical, pharmacy and patient/PPI review pending